INSECT	ICIDAL PROTEINS FROM BACILLUS THURINGIENSIS
Inventors: Gr Jeroen Van R	
	ventis CropScience N.V.
Priority claim	ed of: US patent application No. 60/173387, filed December 28, 1999

•

INSECTICIDAL PROTEINS FROM BACILLUS THURINGIENSIS

INTRODUCTION

The present invention relates to new DNA sequences encoding insecticidal proteins produced by *Bacillus thuringiensis* strains. Particularly, new DNA sequences encoding proteins designated as Cry9Fa, Cry1Jd, and Cry1Bf are provided which are useful to protect plants from insect damage. Also included herein are micro-organisms and plants transformed with at least one of the newly isolated genes so that they are useful to confer insect resistance by expression of insecticidal protein.

BACKGROUND OF THE INVENTION

(i) Field of the Invention:

Bt or *Bacillus thuringiensis* is well known for its specific toxicity to insect pests, and has been used since almost a century to control insect pests of plants. In more recent years, transgenic plants expressing *Bt* proteins were made which were found to successfully control insect damage on plants (e.g., Vaeck et al., 1987).

Despite the isolation of a number of *Bt* crystal protein genes, the search for new genes encoding insecticidal proteins continues. Indeed, insecticidal *Bt* crystal proteins are known to have a relatively narrow target insect range compared to chemical insecticides. Also, having multiple toxins active on the same target insect species allows the use of proteins having different modes of action so that insect resistance development can be prevented or delayed.

(ii) Description of Related Art:

Previously, several types of Cry1B-, Cry1J-, and Cry9-proteins were identified (see Crickmore et al., 1998, incorporated herein by reference, for all details).

The new Cry1Bf protein has the closest sequence identity with the Cry1Be protein (Payne et al, 1998, US Patent 5,723,758), but still differs in about 14 percent of the amino acid sequence of its toxic protein fragment with the toxic fragment of the Cry1Be protein.

The closest sequence identity with the Cry1Jd toxic fragment was found in the toxic fragment of the Cry1Jc1 protein (US Patent 5,723,758), but the toxic fragments of both proteins still differ in about 18 % of their amino acid sequence.

The closest sequence identity with the Cry9Fa toxic fragment was found with the toxic fragment of the Cry9Ea1protein as described by Midoh et al. (PCT Patent publication WO 98/26073) and Narva et al. (PCT patent publication WO 98/00546), but the toxic fragments of the Cry9Fa and Cry9Ea proteins still differ in about 21 % of their amino acid sequence.

SUMMARY OF THE INVENTION

In accordance with this invention is provided a DNA sequence encoding a protein comprising the amino acid sequence selected from the group consisting of: a) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry1Bf* gene deposited at the BCCM-LMBP under accession number LMBP 3986, b) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry1Jd* gene deposited at the BCCM-LMBP under accession number LMBP 3983, and c) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry9Fa* gene deposited at the BCCM-LMBP under accession number LMBP 3984.

Particularly preferred in accordance with this invention is a DNA sequence encoding a protein comprising the amino acid sequence selected from the group consisting of: the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 2, the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 4, and the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 6; alternatively, a DNA encoding a protein comprising the amino acid sequence of the group selected from: the amino acid sequence of SEQ ID No. 2, the amino acid sequence of SEQ ID No. 4, the amino acid sequence of SEQ ID No. 6; or a DNA sequence comprising the DNA sequence of SEQ ID No. 1, SEQ ID No. 3, or SEQ ID No. 5..

Further, in accordance with this invention are provided DNA sequences encoding at least the following portions of the newly-isolated proteins: the amino acid sequence of SEQ ID No. 2 from amino acid position 1 to amino acid position 640, the amino acid sequence of SEQ ID No. 4 from amino acid position 1 to amino acid position 596, and the amino acid sequence of SEQ ID No. 6 from amino acid position 1 to amino acid position 652.

Further, in accordance with this invention are provided the above DNA sequences comprising an artificial DNA sequence having a different codon usage compared to the naturally occurring DNA sequence but encoding the same protein or its insecticidal fragment.

Even further provided in accordance with this invention is a protein comprising the amino acid sequence selected from the group consisting of: a) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry1Bf* gene deposited at the BCCM-LMBP under accession number LMBP 3986, b) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry1Jd* gene deposited at the BCCM-LMBP under accession number LMBP 3983, and c) the amino acid sequence of the insecticidal trypsin-digestion fragment of the protein encoded by the *cry9Fa* gene deposited at the BCCM-LMBP under accession number LMBP 3984.

Particularly preferred herein is a protein comprising the amino acid sequence selected from the group consisting of: the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 2, the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 4, and the amino acid sequence of an insecticidal fragment of the protein of SEQ ID No. 6; alternatively a protein, comprising the amino acid sequence selected from the group consisting of: the amino acid sequence of SEQ ID No. 2 from amino acid position 1 to amino acid position 640, the amino acid sequence of SEQ ID No. 4 from amino acid position 1 to amino acid position 596, and the amino acid sequence of SEQ ID No. 6 from amino acid position 1 to amino acid position 652; or a protein comprising the amino acid sequence of SEQ ID No. 2, SEQ ID No. 4, or SEQ ID No. 6.

Also provided herein are chimeric genes comprising the DNA as defined above under the control of a plant-expressible promoter, and plant cells, plants or seeds transformed to contain those chimeric genes, particularly plant cells, plants, or seeds selected from the group consisting of: corn, cotton, rice, oilseed rape, Brassica species, eggplant, soybean, potato, sunflower, tomato, sugarcane, tea, beans, tobacco, strawberry, clover, cucumber, watermelon, pepper, oat, barley, wheat, dahlia, gladiolus, chrysanthemum, sugarbeet, sorghum, alfalfa, and peanut. In accordance with this invention, the chimeric gene can be integrated in the nuclear or chloroplast DNA of the plant cells.

Further in accordance with this invention are provided micro-organisms, transformed to contain any of the above DNA sequences, particularly those selected from the genus *Agrobacterium*, *Escherichia*, or *Bacillus*.

Also provided herein is a process for controlling insects, comprising expressing any of the above DNA sequences in a host cell, particularly plant cells, and contacting insects with said host cells, and a process for rendering a plant resistant to insects, comprising transforming plants cells with any of the above DNA sequences or chimeric genes, and regenerating transformed plants from such cells which are resistant to insects.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

In accordance with this invention, DNA sequences encoding new *Bt* toxins have been isolated and characterized. The new genes were designated *cry1Bf*, *cry1Jd* and *cry9Fa*, and their encoded proteins Cry1Bf, Cry1Jd and Cry9Fa.

In accordance with this invention "Cry1Bf protein" refers to any protein comprising the smallest protein fragment of the amino acid sequence of SEQ ID No. 2 which retains insecticidal activity, particularly any protein comprising the amino acid sequence from the amino acid at position 1 to the amino acid at position 640 in SEQ ID No. 2, including but not limited to the complete protein with the amino acid sequence of SEQ ID No. 2. This includes

hybrids or chimeric proteins comprising the smallest toxic protein fragment, as well as proteins containing at least one of the three functional domains of the toxic fragment of SEQ ID No. 2. The term "DNA/protein comprising the sequence X", as used herein, refers to a DNA or protein including or containing at least the sequence X, so that other nucleotide or amino acid sequences can be included at the 5' (or N-terminal) and/or 3' (or C-terminal) end, e.g. (the nucleotide sequence of) a selectable marker protein as disclosed in EP 0 193 259.

In accordance with this invention, "Cry9Fa protein" or "Cry9F protein" refers to any protein comprising the smallest protein fragment of the amino acid sequence of SEQ ID No. 6 which retains insecticidal activity, particularly any protein comprising the amino acid sequence from the amino acid at position 1 to the amino acid at position 652 in SEQ ID No. 6, including but not limited to the complete protein with the amino acid sequence of SEQ ID No. 6. This includes hybrids or chimeric proteins comprising the smallest toxic protein fragment, as well as proteins containing at least one of the three functional domains of the toxic fragment of SEQ ID No. 6.

In accordance with this invention, "Cry1Jd protein" refers to any protein comprising the smallest protein fragment of the amino acid sequence of SEQ ID No. 4 which retains insecticidal activity, particularly any protein comprising the amino acid sequence from the amino acid at position 1 to the amino acid at position 596 in SEQ ID No. 4, including but not limited to the complete protein with the amino acid sequence of SEQ ID No. 4. This includes hybrids or chimeric proteins comprising the smallest toxic protein fragment, as well as proteins containing at least one of the three functional domains of the toxic fragment of SEQ ID No. 4.

As used herein, the terms "cry1Bf DNA", "cry9Fa DNA", or "cry1Jd DNA", refer to any DNA sequence encoding the Cry1Bf, Cry9Fa, or Cry1Jd protein, respectively, as defined above. This includes naturally occurring, artificial or synthetic DNA sequences encoding the newly isolated proteins or their insecticidal fragments as defined above. Also included herein are DNA sequences encoding insecticidal proteins which are similar enough to the coding regions of the genomic DNA sequences deposited or the sequences provided in the sequence listing so that they can (i.e., have the ability to) hybridize to these DNA sequences under stringent

hybridization conditions. Stringent hybridization conditions, as used herein, refers particularly to the following conditions: immobilizing the relevant genomic DNA sequences on a filter, and prehybridizing the filters for either 1 to 2 hours in 50 % formamide, 5 % SSPE, 2x Denhardt's reagent and 0.1 % SDS at 42 ° C or 1 to 2 hours in 6x SSC, 2xDenhardt's reagent and 0.1 % SDS at 68 °C. The denatured labeled probe is then added directly to the prehybridization fluid and incubation is carried out for 16 to 24 hours at the appropriate temperature mentioned above. After incubation, the filters are then washed for 20 minutes at room temperature in 1x SSC, 0.1 % SDS, followed by three washes of 20 minutes each at 68 $^{\circ}$ C in 0.2 x SSC and 0.1 % SDS. An autoradiograph is established by exposing the filters for 24 to 48 hours to X-ray film (Kodak XAR-2 or equivalent) at -70 °C with an intensifying screen. Of course, equivalent conditions and parameters can be used in this process while still retaining the desired stringent hybridization conditions. One of such equivalent conditions includes: immobilizing the relevant genomic DNA sequences on a filter, and prehybridizing the filters for either 1 to 2 hours in 50 % formamide, 5 % SSPE, 2x Denhardt's reagent and 0.1 % SDS at 42 ° C or 1 to 2 hours in 6x SSC, 2xDenhardt's reagent and 0.1 % SDS at 68 °C. The denatured (dig- or radio-)labeled probe is then added directly to the prehybridization fluid and incubation is carried out for 16 to 24 hours at the appropriate temperature mentioned above. After incubation, the filters are then washed for 30 minutes at room temperature in 2x SSC, 0.1 % SDS, followed by 2 washes of 30 minutes each at 68 °C in 0.5 x SSC and 0.1 % SDS. An autoradiograph is established by exposing the filters for 24 to 48 hours to X-ray film (Kodak XAR-2 or equivalent) at -70 °C with an intensifying screen

"Insecticidal activity" of a protein, as used herein, means the capacity of a protein to kill insects when such protein is fed to insects, preferably by expression in a recombinant host such as a plant. "Insect-controlling amounts" of a protein, as used herein, refers to an amount of protein which is sufficient to limit damage on a plant by insects feeding on such plant to commercially acceptable levels, e.g. by killing the insects or by inhibiting the insect development or growth in such a manner that they provide less damage to a plant and plant yield is not significantly adversely affected.

In accordance with this invention, insects susceptible to the new Cry proteins of the invention are contacted with this protein in insect-controlling amounts, preferably insecticidal amounts.

"Cry protein" or "Cry protein of this invention", as used herein, refers to any one of the new proteins isolated in accordance with this invention and identified herein as Cry1Bf, Cry9Fa, or Cry1Jd protein. A Cry protein, as used herein, can be a protein in the full length size, also named a protoxin, or can be in a slightly or fully (e.g., N- and C-terminal truncation) truncated form as long as the insecticidal activity is retained, or can be a combination of different proteins or protein parts in a hybrid or fusion protein. A "Cry protoxin" refers to the full length crystal protein as it is encoded by the naturally-occurring Bt DNA sequence, a "Cry toxin" refers to an insecticidal fragment thereof, particularly the smallest toxic fragment thereof, typically in the molecular weight range of about 60 to about 80 kD as determined by SDS-PAGE electrophoresis. A "cry gene" or "cry DNA", as used herein, is a DNA sequence encoding a Cry protein in accordance with this invention, referring to any of the cry1Bf, cry9Fa, and cry1Jd DNA sequences defined above.

The "smallest toxic fragment" of a Cry protein, as used herein, is that fragment as can be obtained by trypsin or chymotrypsin digestion of the full length solubilized crystal protein that retains toxicity, or that toxic protein fragment encoded by DNA fragments of the Cry protein. This protein will mostly have a short N-terminal and a long C-terminal truncation compared to the protoxin. Although for recombinant expression, toxic fragments starting at or around original amino acid position 1 are a more preferred embodiment in accordance with this invention, it should be noted that besides a C-terminal truncation, some N-terminal amino acids can also be deleted while retaining the insecticidal character of the protein. The N-terminal end of the smallest toxic fragment is conveniently determined by N-terminal amino acid sequence determination of trypsin- or chymotrypsin-treated soluble crystal protein by techniques routinely available in the art.

Dna encoding the Cry proteins of this invention can be isolated in a conventional manner from the *E. coli* strains, deposited on November 25, 1999 at the BCCM-LMBP under accession numbers LMBP 3983, LMBP 3984, LMBP 3985 and LMBP 3986. The encoded Cry proteins can be used to prepare specific monoclonal or polyclonal antibodies in a conventional manner (Höfte et al., 1988). The toxin forms can be obtained by protease (e.g., trypsin) digestion of the Cry protoxins.

The DNA sequences encoding the Cry proteins can be isolated in a conventional manner from the respective strains or can be synthesized based on the encoded amino acid sequence.

The DNA sequences encoding the Cry proteins of the invention were identified by digesting total DNA from isolated *Bt* strains with restriction enzymes; size fractionating the DNA fragments, so produced, into DNA fractions of 5 to 10 Kb; ligating these fractions to cloning vectors; screening the *E. coli*, transformed with the cloning vectors, with a DNA probe that was constructed from a region of known *Bt* crystal protein genes or with a DNA probe based on specific PCR fragments generated from *Bt* DNA using primers corresponding to known *Bt* crystal protein genes.

Also, DNA sequences for use in this invention can be made synthetically. Indeed, because of the degeneracy of the genetic code, some amino acid codons can be replaced with others without changing the amino acid sequence of the protein. Furthermore, some amino acids can be substituted by other equivalent amino acids without significantly changing the insecticidal activity of the protein. Also, changes in amino acid sequence or composition in regions of the molecule, different from those responsible for binding and toxicity (e.g., pore formation) are less likely to cause a difference in insecticidal activity of the protein. Such equivalents of the gene include DNA sequences hybridizing to the DNA sequence of the Cry toxins or protoxins of SEQ ID. No. 2, 4, or 6 under stringent conditions and encoding a protein with the same insecticidal characteristics as the (pro)toxin of this invention, or DNA sequences encoding proteins with an amino acid sequence identity of at least 85 %, preferably at least 90 %, most preferably at least 95 %, with the protein toxin form (from the N-terminus to 2 amino acids beyond conserved sequence block 5 as defined in Schnepf et al., 1998) or with the protein protoxin form of the Cry1Bf, Cry9FA or Cry1Jd proteins of this invention, as determined using the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA, version 10.0; GCG defaults were used within the GAP program; for the amino acid sequence comparisons, the blosum62 scoring matrix was used).

Of course, any other DNA sequence differing in its codon usage but encoding the same protein or a similar protein with substantially the same insecticidal activity, can be constructed, depending on the particular purpose. It has been described in prokaryotic and eucaryotic expression systems that changing the codon usage to that of the host cell is desired for gene

expression in foreign hosts (Bennetzen & Hall, 1982; Itakura, 1977). Furthermore, *Bt* crystal protein genes are known to have no bias towards eucaryotic codons, and to be very AT-rich (Adang et al., 1985, Schnepf et al., 1985). Codon usage tables are available in the literature (Wada et al., 1990; Murray et al., 1989) and in the major DNA sequence databanks (e.g. EMBL at Heidelberg, Germany). Accordingly, synthetic DNA sequences can be constructed so that the same or substantially the same proteins are produced. It is evident that several DNA sequences can be devised once the amino acid sequence of the Cry proteins of this invention is known. Such other DNA sequences include synthetic or semi-synthetic DNA sequences that have been changed in order to inactivate certain sites in the gene, e.g. by selectively inactivating certain cryptic regulatory or processing elements present in the native sequence as described in PCT publications WO 91/16432 and WO 93/09218, or by adapting the overall codon usage to that of a more related host organism, preferably that of the host organism in which expression is desired. When making such genes, the encoded amino acid sequence should be retained to the maximum extent possible, although truncations or minor replacements or additions of amino acids can be done as long as the toxicity of the protein is not negatively affected.

Small modifications to a DNA sequence such as described above can be routinely made by PCR-mediated mutagenesis (Ho et al., 1989, White et al., 1989).

With the term "substantially the same", when referring to a protein, is meant to include a protein that differs in some amino acids, or has some amino acids added (e.g. a fusion protein, see Vaeck et al., 1987) or deleted (e.g. N- or C-terminal truncation), as long as the protein has no major difference in its insecticidal activity.

The term "functional domain" of a Cry toxin as used herein means any part(s) or domain(s) of the toxin with a specific structure that can be transferred to another (Cry) protein for providing a new hybrid protein with at least one functional characteristic (e.g., the binding and/or toxicity characteristics) of the Cry toxin of the invention (Ge et al., 1991). Such parts can form an essential feature of the hybrid *Bt* protein with the binding and/or toxicity characteristics of the Cry protein of this invention. Such a hybrid protein can have an enlarged host range, an improved toxicity and/or can be used in a strategy to prevent insect resistance development (European Patent Publication ("EP") 408 403; Visser et al., 1993).

The 5 to 10 Kb fragments, prepared from total DNA of the Bt strains of the invention, can be ligated in suitable expression vectors and transformed in E. coli, and the clones can then be

screened by conventional colony immunoprobing methods (French et al., 1986) for expression of the toxin with monoclonal or polyclonal antibodies raised against the Cry proteins, or by hybridization with DNA probes.

Also, the 5 to 10 Kb fragments, prepared from total DNA of the *Bt* strains of the invention or fragments thereof cloned and/or subcloned in E.coli, can be ligated in suitable *Bt* shuttle vectors (Lereclus et al., 1992) and transformed in a crystal minus *Bt*-mutant. The clones are then screened for production of crystals (detected by microscopy) or crystal proteins (detected by SDS-PAGE).

The genes encoding the Cry proteins of this invention can be sequenced in a conventional manner (Maxam and Gilbert, 1980; Sanger, 1977) to obtain the DNA sequence. Sequence comparisons indicated that the genes are different from previously described genes encoding protoxins and toxins with activity against Lepidoptera (Höfte and Whiteley, 1989; Crickmore, et al., 1998); and the December 15, 1999 and October 16, 2000 updates on the *Bt* nomenclature website corresponding to the Crickmore et al. (1998) publication, found at:

http://epunix.biols.susx.ac.uk/Home/Neil_Crickmore/Bt/index.html

An insecticidally effective part of the DNA sequences, encoding an insecticidally effective portion of the newly identified Cry protein protoxin forms, can be made in a conventional manner after sequence analysis of the gene. In such fragments, it is preferred that at least the sequence up to the C-terminal end of conserved sequence block 5 of *Bt* proteins (Hofte & Whiteley, 1989; Schnepf et al., 1998), preferably up to two amino acids C-terminal of the conserved sequence block 5, is retained. The amino acid sequence of the Cry proteins can be determined from the DNA sequence of the isolated DNA sequences. By "an insecticidally effective part" of DNA sequences encoding the Cry protein, also referred to herein as "truncated gene" or "truncated DNA", is meant a DNA sequence encoding a polypeptide which has fewer amino acids than the Cry protein protoxin form but which is insecticidal to insects.

In order to express all or an insecticidally effective part of the DNA sequence encoding a Cry protein of this invention in *E. coli*, in other *Bt* strains and in plants, suitable restriction sites can be introduced, flanking the DNA sequence. This can be done by site-directed mutagenesis, using well-known procedures (Stanssens et al., 1989; White et al., 1989). In order to obtain improved expression in plants, the codon usage of the *cry* gene or insecticidally effective *cry* gene part of this invention can be modified to form an equivalent, modified or artificial gene or gene

part in accordance with PCT publications WO 91/16432 and WO 93/09218; EP 0 358 962 and EP 0 359 472, or the *Bt* genes or gene parts can be inserted in the chloroplast genome and expressed there using a chloropast-active promoter (e.g., Mc Bride et al., 1995). For obtaining enhanced expression in monocot plants such as corn, a monocot intron also can be added to the chimeric gene, and the DNA sequence of the *cry* gene or its insecticidal part of this invention can be further changed in a translationally neutral manner, to modify possibly inhibiting DNA sequences present in the gene part by means of site-directed intron insertion and/or by introducing changes to the codon usage, e.g., adapting the codon usage to that most preferred by the specific plant (Murray et al., 1989) without changing significantly the encoded amino acid sequence.

Furthermore, the binding properties of the Cry proteins of the invention can be evaluated, using methods known in the art (Van Rie et al., 1990), to determine if the Cry proteins of the invention bind to sites on the insect midgut that are different from those recognized by other, known Cry or other *Bt* proteins. *Bt* toxins with different binding sites in relevant susceptible insects are very valuable to replace known *Bt* toxins to which insects may have developed resistance, or to use in combination with *Bt* toxins having a different mode of action to prevent or delay the development of insect resistance against *Bt* toxins, particularly when expressed in a plant. Because of the characteristics of the newly isolated *Bt* toxins, they are extremely useful for transforming plants, e.g. monocots such as corn or rice and vegetables such as *Brassica* species plants, to protect these plants from insect damage.

The insecticidally effective *cry* gene part or its equivalent, preferably the *cry* chimeric gene, encoding an insecticidally effective portion of the Cry protoxin, can be stably inserted in a conventional manner into the nuclear genome of a single plant cell, and the so-transformed plant cell can be used in a conventional manner to produce a transformed plant that is insect-resistant. In this regard, a disarmed Ti-plasmid, containing the insecticidally effective *cry* gene part, in *Agrobacterium tumefaciens* can be used to transform the plant cell, and thereafter, a transformed plant can be regenerated from the transformed plant cell using the procedures described, for example, in EP 0 116 718, EP 0 270 822, PCT publication WO 84/02913 and published European Patent application ("EP") 0 242 246 and in Gould et al. (1991). Preferred Ti-plasmid vectors each contain the insecticidally effective *cry* gene part between the border sequences, or at least located to the left of the right border sequence, of the T-DNA of the Ti-

plasmid. Of course, other types of vectors can be used to transform the plant cell, using procedures such as direct gene transfer (as described, for example in EP 0 233 247), pollen mediated transformation (as described, for example in EP 0 270 356, PCT publication WO 85/01856, and US Patent 4,684,611), plant RNA virus-mediated transformation (as described, for example in EP 0 067 553 and US Patent 4,407,956), liposome-mediated transformation (as described, for example in US Patent 4,536,475), and other methods such as the recently described methods for transforming certain lines of corn (Fromm et al., 1990; Gordon-Kamm et al., 1990) and rice (Shimamoto et al., 1989; Datta et al., 1990) and the recently described method for transforming monocots generally (PCT publication WO 92/09696).

The resulting transformed plant can be used in a conventional plant breeding scheme to produce more transformed plants with the same characteristics or to introduce the insecticidally effective *cry* gene part in other varieties of the same or related plant species. Seeds, which are obtained from the transformed plants, contain the insecticidally effective *cry* gene part as a stable genomic insert. Cells of the transformed plant can be cultured in a conventional manner to produce the insecticidally effective portion of the Cry protoxin, preferably the Cry toxin, which can be recovered for use in conventional insecticide compositions against Lepidoptera (US Patent 5,254,799). In accordance with this invention, plants or seeds of the invention can be used to obtain resistance to insects, e.g. by sowing or planting in a field wherein damaging insects usually occur, said seeds or plants. Methods for obtaining insect resistance and methods for obtaining improved yield or reduced insect damage are thus provided in accordance with the invention by planting or sowing in a field, preferably a field wherein damaging insects feeding on such plants usually occur or are expected to occur at levels which provide economic damage to the plants, the plants of seeds of the invention producing the Cry proteins of the invention.

The insecticidally effective *cry* gene part, preferably the truncated *cry* gene, is inserted in a plant cell genome so that the inserted gene is downstream (i.e., 3') of, and under the control of, a promoter which can direct the expression of the gene part in the plant cell. This is preferably accomplished by inserting the *cry* chimeric gene in the plant cell genome, particularly in the nuclear or chloroplast genome. Preferred promoters include: the strong constitutive 35S promoters (the "35S promoters") of the cauliflower mosaic virus (CaMV) of isolates CM 1841 (Gardner et al., 1981), CabbB-S (Franck et al., 1980) and CabbB-JI (Hull and Howell, 1987); promoters from the ubiquitin family (e.g., the maize ubiquitin promoter of Christensen

et al., 1992, see also Cornejo et al., 1993), the gos2 promoter (de Pater et al., 1992), the emu promoter (Last et al., 1990), rice actin promoters such as the promoter described by Zhang et al. (1991); and the TR1' promoter and the TR2' promoter (the "TR1' promoter" and "TR2' promoter", respectively) which drive the expression of the 1' and 2' genes, respectively, of the T-DNA (Velten et al., 1984). Alternatively, a promoter can be utilized which is not constitutive but rather is specific for one or more tissues or organs of the plant (e.g., leaves and/or roots) whereby the inserted *cry* gene part is expressed only in cells of the specific tissue(s) or organ(s). For example, the insecticidally effective *cry* gene part could be selectively expressed in the leaves of a plant (e.g., corn, cotton) by placing the insecticidally effective gene part under the control of a light-inducible promoter such as the promoter of the ribulose-1,5-bisphosphate carboxylase small subunit gene of the plant itself or of another plant such as pea as disclosed in US Patent 5,254,799. Another alternative is to use a promoter whose expression is inducible (e.g., by temperature, wounding or chemical factors).

The insecticidally effective *cry* gene part is inserted in the plant genome so that the inserted gene part is upstream (i.e., 5') of suitable 3' end transcription regulation signals (i.e., transcript formation and polyadenylation signals). This is preferably accomplished by inserting the *cry* chimeric gene in the plant cell genome. Preferred polyadenylation and transcript formation signals include those of the octopine synthase gene (Gielen et al., 1984) and the T-DNA gene 7 (Velten and Schell, 1985), which act as 3'-untranslated DNA sequences in transformed plant cells.

The insecticidally effective *cry* gene part can optionally be inserted in the plant genome as a hybrid gene (US Patent 5,254,799; Vaeck et al., 1987) under the control of the same promoter as a selectable marker gene, such as the *neo* gene (EP 0 242 236) encoding kanamycin resistance, so that the plant expresses a fusion protein.

All or part of the *cry* gene, encoding an anti-lepidopteran protein, can also be used to transform other bacteria, such as a *B. thuringiensis* which has insecticidal activity against Lepidoptera or Coleoptera. Thereby, a transformed *Bt* strain can be produced which is useful for combatting a wide spectrum of lepidopteran and coleopteran insect pests or for combatting additional lepidopteran insect pests. Transformation of bacteria, such as bacteria of the genus *Agrobacterium*, *Bacillus* or *Escherichia*, with all or part of the *cry* gene of this invention, incorporated in a suitable cloning vehicle, can be carried out in a conventional manner, preferably

using conventional electroporation techniques as described in Mahillon et al. (1989) and in PCT Patent publication WO 90/06999.

Transformed *Bacillus* species strains containing the *cry* gene of this invention can be fermented by conventional methods (Dulmage, 1981; Bernhard and Utz, 1993) to provide high yields of cells. Under appropriate conditions which are well understood (Dulmage, 1981), these strains each sporulate to produce crystal proteins containing the Cry protoxin in high yields.

An insecticidal, particularly anti-lepidopteran, composition of this invention can be formulated in a conventional manner using the microorganisms transformed with the *cry* gene, or preferably their respective Cry proteins or the Cry protoxin, toxin or insecticidally effective protoxin portion as an active ingredient, together with suitable carriers, diluents, emulsifiers and/or dispersants (e.g., as described by Bernhard and Utz, 1993). This insecticide composition can be formulated as a wettable powder, pellets, granules or dust or as a liquid formulation with aqueous or non-aqueous solvents as a foam, gel, suspension, concentrate, etc.

A method for controlling insects, particularly Lepidoptera, in accordance with this invention can comprise applying (e.g., spraying), to a locus (area) to be protected, an insecticidal amount of the Cry proteins or host cells transformed with the *cry* gene of this invention. The locus to be protected can include, for example, the habitat of the insect pests or growing vegetation or an area where vegetation is to be grown.

To obtain the Cry protoxin or toxin, cells of the recombinant hosts expressing the Cry protein can be grown in a conventional manner on a suitable culture medium and then lysed using conventional means such as enzymatic degradation or detergents or the like. The protoxin can then be separated and purified by standard techniques such as chromatography, extraction, electrophoresis, or the like. The toxin can then be obtained by trypsin digestion of the protoxin.

The following Examples illustrate the invention, and are not provided to limit the invention or the protection sought. The sequence listing referred to in the Examples, the Claims and the Description is as follows:

Sequence Listing:

SEQ ID No. 1 - amino acid and DNA sequence of Cry1Bf protein and DNA

SEQ ID No. 2 - amino acid sequence of Cry1Bf protein.

SEQ ID No. 3 - amino acid and DNA sequence of Cry1Jd protein and DNA.

SEQ ID No. 4 - amino acid sequence Cry1Jd protein.

SEQ ID No. 5 - amino acid and DNA sequence of Cry9Fa protein and DNA.

SEQ ID No. 6 - amino acid sequence of Cry9Fa protein.

SEQ ID No. 7 - DNA sequence for primer Cry1B.fw.

SEQ ID No. 8 - DNA sequence for primer B.R.

SEQ ID No. 9 - DNA sequence for primer B.F.

SEQ ID No. 10 - DNA sequence for primer JFW.

SEQ ID No. 11 - DNA sequence for primer JRV.

SEQ ID No. 12- DNA sequence for primer 9FW.

SEQ ID No. 13 - DNA sequence for primer 9RV.

Unless otherwise stated in the Examples, all procedures for making and manipulating recombinant DNA are carried out by the standard procedures described in Sambrook et al., Molecular Cloning - A Laboratory Manual, Second Ed., Cold Spring Harbor Laboratory Press, NY (1989), and in Volumes 1 and 2 of Ausubel et al. (1994) Current Protocols in Molecular Biology, Current Protocols, USA. Standard materials and methods for plant molecular biology work are described in Plant Molecular Biology Labfax (1993) by R.R.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications (UK). Procedures for PCR technology can be found in "PCR protocols: a guide to methods and applications", Edited by M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White (Academic Press, Inc., 1990).

EXAMPLES

Example 1: Characterization of the strains.

The BtS02072BG strain was isolated from a grain dust sample collected in Santo Tomas la Union, Ilocos, Philippines. The BtS02739C strain was isolated from a grain dust sample collected in Lucena City, South Tagalog, Philippines.

Each strain can be cultivated on conventional standard media, preferably T_3 medium (tryptone 3 g/l, tryptose 2 g/l, yeast extract 1.5 g/l, 5 mg MnCl₂, 0.05 M Na₂HPO₄.2H₂O, 0.05 M

 $NaH_2PO_4.H_2O$, pH 6.8 and 1.5% agar), preferably at 28 °C. For long term storage, it is preferred to mix an equal volume of a spore-crystal suspension with an equal volume of 50% glycerol and store this at -70 °C or lyophilize a spore-crystal suspension. For sporulation, growth on T_3 medium is preferred for 72 hours at 28 °C, followed by storage at 4 °C. The crystal proteins produced by the strains during sporulation are packaged in crystals.

Example 2: Insecticidal activity of the BtS02072BG and BtS02739C strains against selected lepidopteran insect species.

Toxicity assays were performed on neonate larvae of Helicoverpa zea, Heliothis virescens, Ostrinia nubilalis, Spodoptera frugiperda and Sesamia nonagrioides fed on an artificial diet layered with spore-crystal mixtures from either BtS02072BG or BtS02739C, at about 10⁹ spore-crystals per ml.

The artificial diet (Vanderzant, 1962) was dispensed in wells of Costar 24-well plates for tests on H. zea, H. virescens and O. nubilalis. 50 microliter of the spore-crystal mixture was applied on the surface of the diet and dried in a laminar air flow. For tests on H. zea, H. virescens, one larva was placed in each well and 20 larvae were used per sample. For tests on O. nubilalis, 2 larvae were placed in each well and 24 larvae were used per sample. The artificial diet was dispensed in wells of Costar 48-well plates for tests on S. frugiperda and S. nonagrioides. 25 microliter of the spore-crystal mixture was applied on the surface of the diet and dried in a laminar air flow. One larva was placed in each well and 18 larvae were used per sample. Dead and living larvae were counted on the seventh day. The percentage of dead larvae are shown in Table I below.

Table 1: Percentage of dead larvae upon application of crystal-spore mixture to insects:

	BTS02072BG	BTS02739C
H. zea	70	15
H. virescens	85-50	80-60
O. nubilalis	92	72
S. frugiperda	6	Not tested
S. nonagroides	100	Not tested

Example 3: Characterization of new cry genes

The BtS02739C genes were detected by PCR using degenerate primers targeting conserved regions in known *cry* genes. The resulting amplification product was purified using the Wizard PCR preps (Promega) purification system and ligated into pGEM-T vector (Promega). The ligation mixture was electroporated into *E. coli* JM101. A miniprep was made of at least 40 insert-containing transformants, and digests were performed with selected restriction enzymes. Following electrophoresis of the digested miniprep DNA, different DNA fragment patterns could be observed. For each pattern at least one colony was selected. An appropriate DNA prep was made in order to determine the sequence of the insert of the plasmid present in each selected colony. Alignment of the determined sequences of the amplification products with publicly available *cry* sequences demonstrates that strain BtS02739C contains a novel *cry1J*- type gene and a novel *cry9*- type gene.

The BtS02072BG gene was detected as follow. First, a PCR was performed using degenerate crystal protein gene primers on strain BtS02419J. The resulting amplification product was used as template in a secondary PCR using degenerate crystal protein primers

The resulting amplification product was purified using the Wizard PCR preps (Promega) purification system and ligated into pGEM-T vector (Promega). The ligation mixture was electroporated into XL1 Blue *E. coli*. A miniprep was made of at least 40 insert-containing transformants, and digests were performed with selected restriction enzymes. Following electrophoresis of the digested miniprep DNA, different DNA fragment patterns could be observed. For each pattern at least one colony was selected. An appropriate DNA prep was made in order to determine the sequence of the insert of the plasmid present in each selected colony.

From the cloned amplification products from strain BtS02419J, a sequence was found to be identical to the corresponding fragment of *cry1Be1*, except for one nucleotide difference. Next, primers were selected to evaluate the presence of a *cry* sequence similar to that of the sequenced *cry* gene fragment from BtS02419J in a number of Bt strains, one of them being strain BtS02072BG. These primers had the following sequence (5' to 3'):

Forward primer: cry1B.fw: CAG TCC AAA CGG GTA TAA AC

Reverse primer: B.R:

CTG CTT CGA AGG TTG CAG TA

Alignment of the determined sequences from the amplification products with publicly available *cry* sequences demonstrates that strain BtS02072BG contains a novel *cry1B*-type gene.

Example 4: Cloning and expression of the cry genes

In order to isolate the full length *cry1J*- type and *cry9*- type gene from BtS02739C, and the *cry1B*- type gene from BtS02072BG, total DNA from these strains was prepared and partially digested with Sau3A. The digested DNA was size fractionated on a sucrose gradient and fragments ranging from 5 Kb to 10 Kb were ligated to the BamH1-digested and TsAP (thermosensitive alkaline phosphatase)- treated cloning vector pUC19 (Yannisch-Perron et al, 1985). The ligation mixture was electroporated in *E. coli* XL1-Blue or *E. coli* JM109 cells. Transformants were plated on LB-triacillin plates containing Xgal and IPTG and white colonies were selected to be used in filter hybridization experiments. Recombinant *E. coli* clones containing the vector were then screened with the appropriate DIG labeled probes. These probes were prepared as follows. First, a PCR was performed using as template cells from a recombinant *E. coli* clone containing a plasmid harboring the particular *cry* gene fragment, previously amplified using appropriate primers as shown in Table II.

Table II: primers used to isolate novel Bt DNA sequences (Y = C or T, S = G or C):

strain	gene	primer	Length of	Primer sequence
			amplified	
			fragment	
2739C	cry1J-type	JFW	365 bp	GCA GCT AAT GCT ACC ACA TC
	1 57 17 17	JRV		GTG GCG GTA TGC TGA CTA AT
	cry9-type	9FW	576	GYT TTT ATT CGC CCG CCA CA
	1 3.75 37.83	9RV		CGA CAG TAG SAC CCA CTA CT
2072BG	cry1B-type	B.F	922	CAG CGT ATT AAG TCG ATG GA
	1	B.R		CTG CTT CGA AGG TTG CAG TA

The resulting amplification product was gel-purified and used as template in a secondary PCR reaction using DIG- labeled dNTPs. An appropriate amount of this amplification product was used in hybridization reactions.

Colony hybridization for strain BtS02739C was performed with a mixture of the cry1J- type probe and the cry9- type probe. Positive colonies were then hybridized with each probe separately. Colony hybridization for strain BtS02072BG was performed with the cry1B- type probe. Following identification of a positive colony containing a plasmid harboring the full length cry gene, the sequence of the cry gene was determined using the dye terminator labeling method and a Perkin Elmer ABI Prism-377 DNA sequencer for both strands. Upon DNA sequencing, the genes were termed as follows: the cry1J- type and cry9- type gene from BtS02739C were named cry1Jd and cry9Fa, respectively, and the cry1B- type gene from BtS02072BG was named cry1Bf. The genomic sequences of the isolated cry1Jd, cry9Fa, and cry1Bf genes, as well as the proteins they encode, are shown in the Sequence Listing included in this application. Comparison of the sequences with known Cry DNA or protein sequences showed that the sequences are novel and differ in a substantial number of nucleotides or amino acids from known Bt genes and proteins. Tables III-V provide an overview of the sequence identity with respect to the coding regions of the most similar genes and proteins (both protoxin as toxin forms) as determined using the GAP program of the Wisconsin package of GCG (Madison, Wisconsin, USA) version 10.0. GCG defaults were used within the GAP program. For nucleic acid sequence comparisons, the nwsgapdna scoring matrix was used, for amino acid sequence comparisons, the blosum62 scoring matrix. The toxin form, as used in Tables III-V, refers to the protein starting at the first amino acid and ending two amino acids beyond the last amino acid (usually a proline) of conserved sequence block 5, as defined in Schnepf et al. (1998). The protoxin form refers to the entire protein or coding region of the Bt protein/gene.

Table III: Sequence identities for cry1Bf/Cry1Bf:

crylBal	cry1Bb1	cry1Bc1	cry1Bd1	cry1Be1
91.912	83.890	77.207	83.565	93.774
86.562	74.922	74.922	75.342	89.220
Cry1Ba1	Cry1Bb1	Cry1Bc1	Cry1Bd1	Cry1Be1
89.869	80.193	75.795	80.933	92.170
82.520	67.868	67.868	70.142	86.499
	91.912 86.562 Cry1Ba1 89.869	91.912 83.890 86.562 74.922 Cry1Ba1 Cry1Bb1 89.869 80.193	91.912 83.890 77.207 86.562 74.922 74.922 Cry1Ba1 Cry1Bb1 Cry1Bc1 89.869 80.193 75.795	91.912 83.890 77.207 83.565 86.562 74.922 74.922 75.342 Cry1Ba1 Cry1Bb1 Cry1Bc1 Cry1Bd1 89.869 80.193 75.795 80.933

Table IV: Sequence identities for cry9Fa/Cry9Fa:

cry9Aa1	cry9Ba1	cry9Ca1	cry9Da1	cry9Ea1
-				
71.592	78.212	76.614	81.197	84.043
51.782	62.720	68.215	75.593	81.618
Cry9Aa1	Cry9Ba1	Cry9Ca1	Cry9Da1	Cry9Ea1
62.445	72.064	71.553	76.963	82.578
35.828	52.167	59.133	68.372	78.858
	71.592 51.782 Cry9Aa1 62.445	71.592 78.212 51.782 62.720 Cry9Aa1 Cry9Ba1 62.445 72.064	71.592 78.212 76.614 51.782 62.720 68.215 Cry9Aa1 Cry9Ba1 Cry9Ca1 62.445 72.064 71.553	71.592 78.212 76.614 81.197 51.782 62.720 68.215 75.593 Cry9Aa1 Cry9Ba1 Cry9Ca1 Cry9Da1 62.445 72.064 71.553 76.963

Table V: Sequence identities for cry1Jd/Cry1Jd:

DNA	crylJal	cry1Jb1	cry1Jc1
protoxin	83.233	83.176	86.323
toxin	79.526	81.162	88.143
protein	Cry1Ja1	Cry1Jb1	Cry1Jc1
protoxin	79.759	78.830	82.489
toxin	71.574	74.746	81.711

Genomic clones of the newly isolated genes have been deposited at the BCCMTM-LMBP (Belgian Coordinated Collections of Microorganisms - Laboratorium voor Moleculaire Biologie-Plasmidencollectie, University of Gent, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium) under the following accession numbers:

- LMBP 3983 for *E coli* JM109 containing plasmid pUC2739C/1Jd1 comprising the *cry1Jd* gene, deposited on November 25, 1999 (this gene can be isolated from this plasmid on an about 8.4 kb DNA fragment by digestion with XhoI and Smal);
- LMBP 3984 for *E coli* JM109 containing plasmid pUC2739C/9Fa1 comprising the *cry9Fa* gene, deposited on November 25, 1999 (this gene can be isolated from this plasmid on an about 8 kb DNA fragment by digestion with SacI and PstI); and
- LMBP 3986 for *E coli* XL1Blue containing plasmid pUC2072BG/1Bf1 comprising the *cry1Bf* gene, deposited on November 25, 1999 (this gene can be isolated from this plasmid on an about 7 kb DNA fragment by digestion with Sacl and Sall).

Example 5: Insecticidal activity of the cry genes:

The insert containing the *cry9Fa* gene was subcloned into a suitable shuttle vector and the resulting plasmid pSL2739C/9Fa1 was introduced by routine procedures into a crystal-minus *Bt* strain. The crystal protein produced by a sporulated culture of this recombinant *Bt* strain was tested on neonate larvae of *H. virescens* and *O. nubilalis* at a concentration of about 10⁹ particles/ml. On *O. nubilalis* larvae, 100% mortality was observed, whereas 72% mortality was observed on *H. virescens* larvae, whereas after treatment with the crystal-minus control strain all larvae survived.

The insert containing the *cry1Bf* gene was subcloned into a suitable shuttle vector and the resulting plasmid pSL2072BG/1Bf was introduced by routine procedures into a crystal-minus *Bt* strain. The crystal protein produced by a sporulated culture of this recombinant *Bt* strain was tested on larvae of *Sesamia nonagrioides*, *Heliothis virescens*, *Helicoverpa zea* and *O. nubilalis* at different concentrations. Significant high mortality of the Cry1Bf toxin was observed on *H. virescens*, *Ostrinia nubilialis* and *Sesamia nonagrioides*, while lower toxicity was found on *Helicoverpa zea*. After treatment with the crystal-minus control strain all larvae survived.

The insert containing the *cry1Jd* gene was subcloned into a suitable shuttle vector and the resulting plasmid pGl2739C/1Jd was introduced by routine procedures into a crystal-minus *Bt* strain. The crystal protein produced by a sporulated culture of this recombinant *Bt* strain is tested on larvae of *Heliothis virescens at* different concentrations, and significant mortality of the Cry1Jd toxin was observed. After treatment with the crystal-minus control strain all larvae survived.

Example 6: production of the novel Cry proteins in transformed plants.

Chimeric genes encoding the truncated forms of the Cry1Bf, Cry1Jd, and Cry9Fa proteins are made as described in EP 0 193 259 and published PCT patent application WO 94/12264, using the CaMV 35S (Hull and Howell, 1987) and ubiquitin (Christensen et al., 1992)

promoters. Preferably, the codon usage of the open reading frame is adapted to that of the host plant so as to optimize expression efficiency, as described in published PCT patent application WO 94/12264.

Rice, cotton and corn cells are transformed with the resulting chimeric genes.

Corn cells are stably transformed by either *Agrobacterium*-mediated transformation (Ishida et al., 1996, and U.S. Patent No. 5,591,616) or by electroporation using wounded and enzymedegraded embryogenic callus, as described in WO 92/09696 or US Patent 5,641,664 (incorporated herein by reference).

Cotton cells are stably transformed by *Agrobacterium*-mediated transformation (Umbeck et al., 1987, Bio/Technology 5, 263-266; US Patent 5,004,863, incorporated herein by reference). Rice cells are stably transformed with the method described in published PCT patent application WO 92/09696.

Regenerated transformed corn, cotton and rice plants are selected by ELISA, Northern and Southern blot and insecticidal effect. Chimeric gene-containing progeny plants show improved resistance to insects compared to untransformed control plants with appropriate segregation of insect resistance and the transformed phenotype. Protein and RNA measurements show that increased insect resistance is linked with higher expression of the novel Cry protein in the plants.

REFERENCES CITED

Adang et al.(1985). Gene 36, 289.

Bennetzen & Hall.(1982).J. Biol. Chem. 257, 3026-3031.

Berhard, K. and Utz, R., "Production of *Bacillus thuringiensis* insecticides for experimental and commercial uses", In *Bacillus thuringiensis*, An Environmental Biopesticide: Theory and Practice, pp.255-267, eds. Entwistle, P.F., Cory, J.S., Bailey, M.J. and Higgs, S., John Wiley and Sons, New York (1993).

Christensen et al. (1992) Plant mol. Biol. 18, 675-689.

Cornejo et al. (1993) Plant Mol. Biol. 23, 567-581.

Crickmore et al. (1998) Microbiol. Mol. Biol Rev. 62(3), 807-13.

Datta S., Peterhans A., Datta K. and Potrykus I., Bio/Technology 8, 736-740 (1990).

Dulmage, H.T., "Production of Bacteria for Biological Control of Insects" in Biological Control in Crop Production, Ed. Paparizas, D.C., Osmun Publishers, Totowa, N.J., USA, pp. 129-141 (1981).

Finney, Probit Analysis, 3rd Edition, Cambridge University Press (1971)

Franck, Guilley, Jonard, Richards and Hirth, Cell 21, 285-294 (1980)

French, B.T., Maul, H.N. and Maul, G.G., Anal.Biochem. 156, 417-423 (1986)

Fromm M., Morrish F., Armstrong C., Williams R., Thomas J. and Klein T., Bio/Technology 8, 833-839 (1990).

Gardner, Howarth, Hahn, Brown-Luedi, Shepard and Messing, Nucleic Acids Research 9, 2871-2887 (1981)

Ge A., Rivers D., Milne R. and Dean D., J. Biol. Chem. 266, 17954-17958 (1991)

Gielen, J., De Beukeleer, M., Seurinck, J., Deboeck, F., De Greve, H., Lemmers, M., Van Montagu, M. and Schell, J., EMBO J 3, 835-845 (1984).

Gordon-Kamm W., Spencer M., Mangano M., Adams T., Daines R., Start W., O'Brien J., Chambers S., Adams W., Willets N., Rice T., Mackey C., Krueger R., Kausch A. and Lemaux P., The Plant Cell 2, 603-618 (1990).

Gould, J., Devey, M., Hasegawa, O., Ulian, E.C., Peterson, G. and Smith, R.H., Plant Physiol. 95, 426-434 (1991).

Ho et al.(1989). Gene 77, 51-59.

Höfte, H., Van Rie, J., Jansens, S., Van Houtven, A., Verbruggen, H. and Vaeck, M., Applied and Environmental Microbiology 54, 2010-2017 (1988)

Höfte H. and Whiteley H.R., Microbiological Review 53, 242-255 (1989).

Hull and Howell, Virology 86, 482-493 (1987)

Ishida et al., 1996, Nature Biotechnology 14, 745-750.

Itakura et al.(1977). Science 198, 1056-1063.

Jansens et al. (1997) Crop Science 37, 1616-1624.

Last et al. (1990) Theor. Appl. Genet. 81, 581-588.

Lereclus, D.; Vallade, M.; Chaufaux, J.; Arantes, O. & Rambaud, S., Bio/Technology 10, 418 (1992).

Mahillon et al, FEMS Microbiol. Letters 60, 205-210 (1989).

Maxam, A.M. and Gilbert, W., Methods in Enzymol. 65, 499-560 (1980).

McBride et al., 1995, Bio/Technology 13, 362

Murray, E., Lotzer, J. and Eberle, M., Nucleic Acids Research 17(2), 477-498 (1989).

Sanger et al., Proc. Natl. Acad. Sci. U S A. 74(12), 5463-5467 (1977).

Schnepf et al.(1985). Journal of Biological Chemistry 260, 6264.

Schnepf et al. (1998). Microbiol. Mol. Biol. Rev. 62(3), 775-806.

Shimamoto K., Terada R., Izawa T. and Fujimoto H., Nature 338, 274-276 (1989).

Stanssens P., Opsomer C., McKeown Y., Kramer W., Zabeau M. and Fritz H.J., Nucleic Acids Research 12, 4441-4454 (1989).

Vaeck, M., Reynaerts, A., Höfte, H., Jansens, S., De Beuckeleer, M., Dean, C., Zabeau, M.,

Vanderzant, J. Econ. Entomol. 55, p. 140 (1962).

Van Montagu, M. and Leemans, J., Nature 327, 33-37(1987).

Van Rie et al., Science 247, 72 (1990).

Velten, J., Velten, L., Hain, R. and Schell, J., EMBO J 3, 2723-2730 (1984).

Velten, J. and Schell, J. Nucleic Acids Research 13, 6981-6998 (1985)

Visser, B., Bosch, D. and Honée, G., "Domain-Structure Studies of *Bacillus thuringiensis* Crystal Proteins: A Genetic Approach", In *Bacillus thuringiensis*, An Environmental Biopesticide: Theory and Practice, pp.71-88, eds. Entwistle, P.F., Cory, J.S., Bailey, M.J. and Higgs, S., John Wiley and Sons, New York (1993).

Wada et al. (1990). Nucl. Acids Res. 18, 2367-1411.

White et al.(1989). Trends in Genet. 5, 185-189.

Yannisch-Perron, C., Vierra, J. and Messing, J., Gene 33, 103-119 (1985).

Zhang et al. (1991) The Plant Cell 3, 1155-1165.

SEQUENCE LISTING

```
<110> Arnaut, Greta
      Boets, Annemie
      Damme, Nicole
      Mathieu, Eva
      Vanneste, Stijn
      Van Rie, Jeroen
<120> Insecticidal proteins from Bacillus thuringiensis.
<130> NEWBTSUS2
<140>
<141>
<150> 60/173387
<151> 1999-12-28
<160> 13
<170> PatentIn Ver. 2.0
<210> 1
<211> 3687
<212> DNA
<213> Bacillus thuringiensis
<220>
<221> CDS
<222> (1)..(3687)
<400> 1
ttg act tca aat agg aaa aat gag aat gaa att ata aat gct tta tcg
                                                                    48
Leu Thr Ser Asn Arg Lys Asn Glu Asn Glu Ile Ile Asn Ala Leu Ser
                                      10
                  5 .
  1
                                                                    96
att cca gct gta tcg aat cat tcc aca caa atg gat cta tca cca gat
Ile Pro Ala Val Ser Asn His Ser Thr Gln Met Asp Leu Ser Pro Asp
                                                      30
                                  25
             20
                                                                    144
gct cgt att gag gat tct ttg tgt ata gcc gag ggg aat aat atc aat
Ala Arg Ile Glu Asp Ser Leu Cys Ile Ala Glu Gly Asn Asn Ile Asn
         35
                              40
                                                  45
cca ctt gtt agc gca tca aca gtc caa acg ggt att aac ata gct ggt
                                                                    192
Pro Leu Val Ser Ala Ser Thr Val Gln Thr Gly Ile Asn Ile Ala Gly
     50
                          55
```

							gta Val									240
							gaa Glu									288
							gtc Val									336
							gca Ala 120									384
							cag Gln									432
							agt Ser									480
							gcg Ala									528
							gta Val									576
							tct Ser 200									624
							tat Tyr									672
							gta Val									720
agc Ser	ttg Leu	aga Arg	ggg Gly	aca Thr 245	aat Asn	gcc Ala	gca Ala	agt Ser	tgg Trp 250	gtg Val	cgt Arg	tat Tyr	aat Asn	caa Gln 255	ttc Phe	768
							gta Val									816

agc Ser	tat Tyr	gac Asp 275	act Thr	cgc Arg	act Thr	tat Tyr	cca Pro 280	ata Ile	aat Asn	acg Thr	agt Ser	gct Ala 285	cag Gln	tta Leu	aca Thr	864
agg Arg	gaa Glu 290	gtt Val	tat Tyr	aca Thr	gac Asp	gca Ala 295	att Ile	gga Gly	gca Ala	aca Thr	300 Gly 999	gta Val	aat Asn	atg Met	gca Ala	912
agt Ser 305	atg Met	aat Asn	tgg Trp	tat Tyr	aat Asn 310	aat Asn	aat Asn	gca Ala	cct Pro	tcg Ser 315	ttt Phe	tcc Ser	gct Ala	ata Ile	gag Glu 320	960
act Thr	gcg Ala	gtt Val	atc Ile	cga Arg 325	agc Ser	ccg Pro	cat His	cta Leu	ctt Leu 330	gat Asp	ttt Phe	cta Leu	gaa Glu	caa Gln 335	ctt Leu	1008
aca Thr	att Ile	ttt Phe	agc Ser 340	act Thr	tca Ser	tca Ser	cga Arg	tgg Trp 345	agt Ser	gct Ala	act Thr	agg Arg	cat His 350	atg Met	act Thr	1056
tac Tyr	tgg Trp	cgg Arg 355	gly aaa	cac His	aca Thr	att Ile	caa Gln 360	tct Ser	cgg Arg	cca Pro	ata Ile	gga Gly 365	ggc	gga Gly	tta Leu	1104
aat Asn	acc Thr 370	tca Ser	acg Thr	cat His	gjå aaa	tct Ser 375	acc Thr	aat Asn	act Thr	tct Ser	att Ile 380	aat Asn	cct Pro	gta Val	aga Arg	1152
tta Leu 385	tca Ser	ttc Phe	ttc Phe	tct Ser	cga Arg 390	gac Asp	gta Val	tat Tyr	tgg Trp	act Thr 395	gaa Glu	tca Ser	tat Tyr	gca Ala	gga Gly 400	1200
gtg Val	ctt Leu	cta Leu	tgg Trp	gga Gly 405	att Ile	tac Tyr	ctt Leu	gaa Glu	cct Pro 410	att Ile	cat His	ggt Gly	gtc Val	cct Pro 415	act Thr	1248
gtt Val	aga Arg	Phe	Asn	ttt Phe	Arg	Asn	Pro	Gln	Asn	act Thr	ttt Phe	gaa Glu	aga Arg 430	ggt Gly	act Thr	1296
gct Ala	aac Asn	tat Tyr 435	agt Ser	caa Gln	ccc Pro	tat Tyr	gag Glu 440	tca Ser	cct Pro	gly aaa	ctt Leu	caa Gln 445	tta Leu	aaa Lys	gat Asp	1344
tca Ser	gaa Glu 450	act Thr	gaa Glu	tta Leu	cca Pro	cca Pro 455	gaa Glu	aca Thr	aca Thr	gaa Glu	cga Arg 460	cca Pro	aat Asn	tat Tyr	gaa Glu	1392
tca Ser 465	Tyr	agt Ser	cat His	agg Arg	tta Leu 470	Ser	cac His	ata Ile	gly ggg	ctc Leu 475	Ile	tca Ser	caa Gln	tct Ser	agg Arg 480	1440

gtg Val	cat His	gta Val	cca Pro	gta Val 485	tat Tyr	tct Ser	tgg Trp	acg Thr	cac His 490	cgt Arg	agt Ser	gca Ala	gat Asp	cgt Arg 495	aca Thr	1488
aat Asn	acc Thr	att Ile	agt Ser 500	tca Ser	gat Asp	agc Ser	ata Ile	aca Thr 505	caa Gln	ata Ile	cca Pro	ttg Leu	gta Val 510	aaa Lys	tca Ser	1536
ttc Phe	aac Asn	ctt Leu 515	aat Asn	tca Ser	ggt Gly	acc Thr	tct Ser 520	gta Val	gtc Val	agt Ser	ggc Gly	cca Pro 525	gga Gly	ttt Phe	aca Thr	1584
gga Gly	999 Gly 530	gat Asp	ata Ile	atc Ile	cga Arg	act Thr 535	aac Asn	gtt Val	aat Asn	ggt Gly	agt Ser 540	gta Val	cta Leu	agt Ser	atg Met	1632
ggt Gly 545	ctt Leu	aat Asn	ttt Phe	aat Asn	aat Asn 550	aca Thr	tca Ser	tta Leu	cag Gln	cgg Arg 555	tat Tyr	cgc Arg	gtg Val	aga Arg	gtt Val 560	1680
cgt Arg	tat Tyr	gct Ala	gct Ala	tct Ser 565	caa Gln	aca Thr	atg Met	gtc Val	ctg Leu 570	agg Arg	gta Val	act Thr	gtc Val	gga Gly 575	Gly 999	1728
agt Ser	act Thr	act Thr	ttt Phe 580	gat Asp	caa Gln	gga Gly	ttc Phe	cct Pro 585	agt Ser	act Thr	atg Met	agt Ser	gca Ala 590	aat Asn	gag Glu	1776
tct Ser	ttg Leu	aca Thr 595	tct Ser	caa Gln	tca Ser	ttt Phe	aga Arg 600	ttt Phe	gca Ala	gaa Glu	ttt Phę	cct Pro 605	gta Val	ggt Gly	att Ile	1824
agt Ser	gca Ala 610	tct Ser	ggc	agt Ser	caa Gln	act Thr 615	gct Ala	gga Gly	ata Ile	agt Ser	ata Ile 620	agt Ser	aat Asn	aat Asn	gca Ala	1872
ggt Gly 625	Arg	Gln	Thr	ttt Phe	His	Phe	gat Asp	Lys	Ile	Glu	Phe	att Ile	cca Pro	att Ile	act Thr 640	1920
gca Ala	acc Thr	ttc Phe	gaa Glu	gca Ala 645	gaa Glu	tac Tyr	gat Asp	tta Leu	gaa Glu 650	agg Arg	gcg Ala	caa Gln	gag Glu	gcg Ala 655	gtg Val	1968
aat Asn	gct Ala	ctg Leu	ttt Phe 660	act Thr	aat Asn	acg Thr	aat Asn	cca Pro 665	aga Arg	aga Arg	ttg Leu	aaa Lys	aca Thr 670	gat Asp	gtg Val	2016
aca Thr	gat Asp	tat Tyr 675	His	att Ile	gat Asp	caa Gln	gta Val 680	tcc Ser	aat Asn	tta Leu	gtg Val	gcg Ala 685	tgt Cys	tta Leu	tcg Ser	2064

gat Asp	gaa Glu 690	ttc Phe	tgc Cys	tta Leu	gat Asp	gaa Glu 695	aag Lys	aga Arg	gaa Glu	tta Leu	ctt Leu 700	gag Glu	aaa Lys	gtg Val	aaa Lys	2112
tat Tyr 705	gcg Ala	aaa Lys	cga Arg	ctc Leu	agt Ser 710	gat Asp	gaa Glu	aga Arg	aac Asn	tta Leu 715	ctc Leu	caa Gln	gat Asp	cca Pro	aac Asn 720	2160
ttc Phe	aca Thr	tcc Ser	atc Ile	aat Asn 725	aag Lys	caa Gln	cca Pro	gac Asp	ttc Phe 730	ata Ile	tct Ser	act Thr	aat Asn	gag Glu 735	caa Gln	2208
tcg Ser	aat Asn	ttc Phe	aca Thr 740	tct Ser	atc Ile	cat His	gaa Glu	caa Gln 745	tct Ser	gaa Glu	cat His	gga Gly	tgg Trp 750	tgg Trp	gga Gly	2256
							gaa Glu 760									2304
tac Tyr	gtc Val 770	aca Thr	cta Leu	ccg Pro	Gly ggg	act Thr 775	ttt Phe	aat Asn	gag Glu	tgt Cys	tat Tyr 780	ccg Pro	acg Thr	tat Tyr	tta Leu	2352
tat Tyr 785	caa Gln	aaa Lys	ata Ile	gga Gly	gag Glu 790	tcg Ser	gaa Glu	tta Leu	aaa Lys	gct Ala 795	tat Tyr	act Thr	cgc Arg	tac Tyr	caa Gln 800	2400
							agt Ser									2448
cgt Arg	tat Tyr 82	Asn	gcg Ala	aaa Lys	cat His	gaa Glu 82	aca Thr 5	ttg Leu	gat Asp	gtt Val	cca Pro 83	Gly	acc Thr	gag Glu	tcc Ser	2496
							agc Ser 840									2544
aat Asn	cga Arg 850	tgc Cys	gca Ala	cca Pro	cat His	ttt Phe 855	gaa Glu	tgg Trp	aat Asn	cct Pro	gat Asp 860	cta Leu	gat Asp	tgt Cys	tcc Ser	2592
tgc Cys 865	aga Arg	gat Asp	gga Gly	gaa Glu	aaa Lys 870	Cys	gcg Ala	cat His	cat His	tcc Ser 875	cat His	cat His	ttc Phe	tct Ser	ttg Leu 880	2640
gat Asp	Ile	gat Asp 885	att Ile	gga Gly	tgc Cys	Thr	gac Asp 890	ttg Leu	cat His	gag Glu	Asn	cta Leu 895	ggc	gtg Val	tgg Trp	2688

gtg Val	gta Val	ttc Phe	aag Lys 900	att Ile	aag Lys	acg Thr	cag Gln	gaa Glu 905	ggt Gly	cat His	gca Ala	aga Arg	cta Leu 910	Gly ggg	aat Asn	2736
ctg Leu	gaa Glu	ttt Phe 915	att Ile	gaa Glu	gag Glu	aaa Lys	cca Pro 920	tta Leu	tta Leu	gga Gly	gaa Glu	gca Ala 925	ctg Leu	tct Ser	cgt Arg	2784
gtg Val	aag Lys 930	aga Arg	gca Ala	gag Glu	aaa Lys	aaa Lys 935	tgg Trp	aga Arg	gac Asp	aaa Lys	cgt Arg 940	gaa Glu	aaa Lys	cta Leu	caa Gln	2832
ttg Leu 945	gaa Glu	aca Thr	aaa Lys	cga Arg	gta Val 950	tat Tyr	aca Thr	gag Glu	gca Ala	aaa Lys 955	gaa Glu	gct Ala	gtg Val	gat Asp	gct Ala 960	2880
	ttt Phe															2928
ggc	atg Met	att Ile	cat His 980	gcg Ala	gca Ala	gat Asp	aaa Lys	ctt Leu 985	gtt Val	cat His	cga Arg	att Ile	cga Arg 990	gag Glu	gct Ala	2976 ,
tat Tyr	ctg Leu	tca Ser 995	gaa Glu	tta Leu	tct Ser	Val	atc Ile 1000	ccg Pro	ggt Gly	gta Val	Asn	gcg Ala 1005	gaa Glu	att Ile	ttt Phe	3024
Glu	gaa Glu 1010	tta Leu	gaa Glu	ggt Gly	Arg	att Ile 1015	atc Ile	act Thr	gca Ala	Ile	tcc Ser 1020	cta Leu	tac Tyr	gat Asp	gcg Ala	3072
aga Arg 102	aat Asn 5	gtc Val	gtt Val	Lys	aat Asn 1030	ggt Gly	gat Asp	ttt Phe	Asn	aat Asn 1035	gga Gly	tta Leu	gca Ala	Cys	tgg Trp 1040	3120
aat Asn	gta Val	aaa Lys	Gly	cat His 1045	Val	gat Asp	Val	Gln	cag Gln 1050	Ser	cat His	cac His	Arg	tct Ser 1055	gtc Val	3168
ctt Leu	gtt Val	Ile	cca Pro 1060	Glu	tgg Trp	gaa Glu	Ala	gaa Glu 1065	gtg Val	tca Ser	caa Gln	Ala	gtt Val 1070	Arg	gtc Val	3216
tgt Cys	Pro	999 Gly 1075	Arg	ggc Gly	tat Tyr	Ile	ctc Leu 1080	Arg	gtc Val	aca Thr	Ala	tac Tyr 1085	Lys	gag Glu	gga Gly	3264
tat Tyr	gga Gly 1090	Glu	ggt Gly	tgt Cys	Val	acg Thr 1095	Ile	cat His	gaa Glu	Ile	gag Glu 1100	Asn	aat Asn	aca Thr	gac Asp	3312

gaa cta aaa ttt aaa aac tgt gaa gaa gag gaa gtg tat cca acg gat Glu Leu Lys Phe Lys Asn Cys Glu Glu Glu Glu Val Tyr Pro Thr Asp 1105 1110 1115 1120	3360
aca gga acg tgt aat gat tat act gca cac caa ggt aca gca gta tgt Thr Gly Thr Cys Asn Asp Tyr Thr Ala His Gln Gly Thr Ala Val Cys 1125 1130 1135	3408
aat tcc cgt aat gct gga tat gag gat gca tat gaa gtt gat act aca Asn Ser Arg Asn Ala Gly Tyr Glu Asp Ala Tyr Glu Val Asp Thr Thr 1140 1145 1150	3456
gca tct gtt aat tac aaa ccg act tat gaa gaa gaa acg tat aca gat Ala Ser Val Asn Tyr Lys Pro Thr Tyr Glu Glu Glu Thr Tyr Thr Asp 1155 1160 1165	3504
gta cga aga gat aat cat tgt gaa tat gac aga ggg tat gtg aat tat Val Arg Arg Asp Asn His Cys Glu Tyr Asp Arg Gly Tyr Val Asn Tyr 1170 1175 1180	3552
cca cca cta cca gct ggt tat atg aca aaa gaa tta gaa tac ttc cca Pro Pro Leu Pro Ala Gly Tyr Met Thr Lys Glu Leu Glu Tyr Phe Pro 1185 1190 1195 1200	3600
gaa acc gat aag gta tgg att gag att gga gaa acg gaa ggg aag ttt Glu Thr Asp Lys Val Trp Ile Glu Ile Gly Glu Thr Glu Gly Lys Phe 1205 1210 1215	3648
att gta gac agc gtg gaa tta ctc ctt atg gag gaa tag Ile Val Asp Ser Val Glu Leu Leu Met Glu Glu 1220 1225	3687
<210> 2 <211> 1228 <212> PRT <213> Bacillus thuringiensis	
<pre><400> 2 Leu Thr Ser Asn Arg Lys Asn Glu Asn Glu Ile Ile Asn Ala Leu Ser 1 5 10 15</pre>	
Ile Pro Ala Val Ser Asn His Ser Thr Gln Met Asp Leu Ser Pro Asp 20 25 30	
Ala Arg Ile Glu Asp Ser Leu Cys Ile Ala Glu Gly Asn Asn Ile Asn	
35 40 45	

- Arg Ile Leu Gly Val Leu Gly Val Pro Phe Ala Gly Gln Ile Ala Ser 65 70 75 80
- Phe Tyr Ser Phe Leu Val Gly Glu Leu Trp Pro Arg Gly Arg Asp Gln 85 90 95
- Trp Glu Ile Phe Leu Glu His Val Glu Gln Leu Ile Asn Gln Gln Ile 100 105 110
- Thr Glu Asn Ala Arg Asn Thr Ala Leu Ala Arg Leu Gln Gly Leu Gly 115 120 125
- Asp Ser Phe Arg Ala Tyr Gln Gln Ser Leu Glu Asp Trp Leu Glu Asn 130 135 140
- Arg Asp Asp Ala Arg Thr Arg Ser Val Leu Tyr Thr Gln Tyr Ile Ala 145 150 155 160
- Leu Glu Leu Asp Phe Leu Asn Ala Met Pro Leu Phe Ala Ile Arg Asn 165 170 175
- Gln Glu Val Pro Leu Leu Met Val Tyr Ala Gln Ala Ala Asn Leu His 180 185 190
- Leu Leu Leu Arg Asp Ala Ser Leu Phe Gly Ser Glu Phe Gly Leu
 195 200 205
- Thr Ser Gln Glu Ile Gln Arg Tyr Tyr Glu Arg Gln Val Glu Gln Thr 210 215 220
- Arg Asp Tyr Ser Asp Tyr Cys Val Glu Trp Tyr Asn Thr Gly Leu Asn 225 230 235 240
- Ser Leu Arg Gly Thr Asn Ala Ala Ser Trp Val Arg Tyr Asn Gln Phe 245 250 255
- Arg Arg Asp Leu Thr Leu Gly Val Leu Asp Leu Val Ala Leu Phe Pro 260 265 270
- Ser Tyr Asp Thr Arg Thr Tyr Pro Ile Asn Thr Ser Ala Gln Leu Thr 275 280 285
- Arg Glu Val Tyr Thr Asp Ala Ile Gly Ala Thr Gly Val Asn Met Ala 290 295 300
- Ser Met Asn Trp Tyr Asn Asn Asn Ala Pro Ser Phe Ser Ala Ile Glu 305 310 315 320
- Thr Ala Val Ile Arg Ser Pro His Leu Leu Asp Phe Leu Glu Gln Leu 325 330 335

- Thr Ile Phe Ser Thr Ser Ser Arg Trp Ser Ala Thr Arg His Met Thr 340 345 350
- Tyr Trp Arg Gly His Thr Ile Gln Ser Arg Pro Ile Gly Gly Leu 355 360 365
- Asn Thr Ser Thr His Gly Ser Thr Asn Thr Ser Ile Asn Pro Val Arg 370 375 380
- Leu Ser Phe Phe Ser Arg Asp Val Tyr Trp Thr Glu Ser Tyr Ala Gly 385 390 395 400
- Val Leu Leu Trp Gly Ile Tyr Leu Glu Pro Ile His Gly Val Pro Thr 405 410 415
- Val Arg Phe Asn Phe Arg Asn Pro Gln Asn Thr Phe Glu Arg Gly Thr
 420 425 430
- Ala Asn Tyr Ser Gln Pro Tyr Glu Ser Pro Gly Leu Gln Leu Lys Asp
 435 440 445
- Ser Glu Thr Glu Leu Pro Pro Glu Thr Thr Glu Arg Pro Asn Tyr Glu 450 455 460
- Ser Tyr Ser His Arg Leu Ser His Ile Gly Leu Ile Ser Gln Ser Arg 465 470 475 480
- Val His Val Pro Val Tyr Ser Trp Thr His Arg Ser Ala Asp Arg Thr 485 490 495
- Asn Thr Ile Ser Ser Asp Ser Ile Thr Gln Ile Pro Leu Val Lys Ser 500 505 510
- Phe Asn Leu Asn Ser Gly Thr Ser Val Val Ser Gly Pro Gly Phe Thr 515 520 525
- Gly Gly Asp Ile Ile Arg Thr Asn Val Asn Gly Ser Val Leu Ser Met 530 535 540
- Gly Leu Asn Phe Asn Asn Thr Ser Leu Gln Arg Tyr Arg Val Arg Val 545 550 555 560
- Arg Tyr Ala Ala Ser Gln Thr Met Val Leu Arg Val Thr Val Gly Gly 565 570 575
- Ser Thr Thr Phe Asp Gln Gly Phe Pro Ser Thr Met Ser Ala Asn Glu 580 585 590
- Ser Leu Thr Ser Gln Ser Phe Arg Phe Ala Glu Phe Pro Val Gly Ile 595 600 605

- Ser Ala Ser Gly Ser Gln Thr Ala Gly Ile Ser Ile Ser Asn Asn Ala 610 615 620
- Gly Arg Gln Thr Phe His Phe Asp Lys Ile Glu Phe Ile Pro Ile Thr 625 630 635 640
- Ala Thr Phe Glu Ala Glu Tyr Asp Leu Glu Arg Ala Gln Glu Ala Val 645 650 655
- Asn Ala Leu Phe Thr Asn Thr Asn Pro Arg Arg Leu Lys Thr Asp Val 660 665 670
- Thr Asp Tyr His Ile Asp Gln Val Ser Asn Leu Val Ala Cys Leu Ser 675 680 685
- Asp Glu Phe Cys Leu Asp Glu Lys Arg Glu Leu Leu Glu Lys Val Lys 690 695 700
- Tyr Ala Lys Arg Leu Ser Asp Glu Arg Asn Leu Leu Gln Asp Pro Asn 705 710 715 720
- Phe Thr Ser Ile Asn Lys Gln Pro Asp Phe Ile Ser Thr Asn Glu Gln 725 730 735
- Ser Asn Phe Thr Ser Ile His Glu Gln Ser Glu His Gly Trp Trp Gly
 740 745 750
- Ser Glu Asn Ile Thr Ile Gln Glu Gly Asn Asp Val Phe Lys Glu Asn 755 760 765
- Tyr Val Thr Leu Pro Gly Thr Phe Asn Glu Cys Tyr Pro Thr Tyr Leu 770 775 780
- Tyr Gln Lys Ile Gly Glu Ser Glu Leu Lys Ala Tyr Thr Arg Tyr Gln 785 790 795 800
- Leu Arg Gly Tyr Ile Glu Asp Ser Gln Asp Leu Glu Ile Tyr Leu Ile 805 810 815
- Arg Tyr Asn Ala Lys His Glu Thr Leu Asp Val Pro Gly Thr Glu Ser 820 825 830
- Val Trp Pro Leu Ser Val Glu Ser Pro Ile Gly Arg Cys Gly Glu Pro 835 840 845
- Asn Arg Cys Ala Pro His Phe Glu Trp Asn Pro Asp Leu Asp Cys Ser 850 855 860
- Cys Arg Asp Gly Glu Lys Cys Ala His His Ser His His Phe Ser Leu 865 870 875 880

- Asp Ile Asp Ile Gly Cys Thr Asp Leu His Glu Asn Leu Gly Val Trp 885 890 895
- Val Val Phe Lys Ile Lys Thr Gln Glu Gly His Ala Arg Leu Gly Asn 900 905 910
- Leu Glu Phe Ile Glu Glu Lys Pro Leu Leu Gly Glu Ala Leu Ser Arg 915 920 925
- Val Lys Arg Ala Glu Lys Lys Trp Arg Asp Lys Arg Glu Lys Leu Gln 930 935 940
- Leu Glu Thr Lys Arg Val Tyr Thr Glu Ala Lys Glu Ala Val Asp Ala 945 950 955 960
- Leu Phe Val Asp Ser Gln Tyr Asn Arg Leu Gln Ala Asp Thr Asn Ile 965 970 975
- Gly Met Ile His Ala Ala Asp Lys Leu Val His Arg Ile Arg Glu Ala 980 985 990
- Tyr Leu Ser Glu Leu Ser Val Ile Pro Gly Val Asn Ala Glu Ile Phe 995 1000 1005
- Glu Glu Leu Glu Gly Arg Ile Ile Thr Ala Ile Ser Leu Tyr Asp Ala 1010 1015 1020
- Arg Asn Val Val Lys Asn Gly Asp Phe Asn Asn Gly Leu Ala Cys Trp 025 1030 1035 1040
- Asn Val Lys Gly His Val Asp Val Gln Gln Ser His His Arg Ser Val 1045 1050 1055
- Leu Val Ile Pro Glu Trp Glu Ala Glu Val Ser Gln Ala Val Arg Val 1060 1065 1070
- Cys Pro Gly Arg Gly Tyr Ile Leu Arg Val Thr Ala Tyr Lys Glu Gly 1075 1080 1085
- Tyr Gly Glu Gly Cys Val Thr Ile His Glu Ile Glu Asn Asn Thr Asp 1090 1095 1100
- Glu Leu Lys Phe Lys Asn Cys Glu Glu Glu Glu Val Tyr Pro Thr Asp 105 1110 1115 1120
- Thr Gly Thr Cys Asn Asp Tyr Thr Ala His Gln Gly Thr Ala Val Cys 1125 1130 1135
- Asn Ser Arg Asn Ala Gly Tyr Glu Asp Ala Tyr Glu Val Asp Thr Thr
 1140 1145 1150
- Ala Ser Val Asn Tyr Lys Pro Thr Tyr Glu Glu Glu Thr Tyr Thr Asp 1155 1160 1165

Val Arg Arg Asp Asn His Cys Glu Tyr Asp Arg Gly Tyr Val Asn Tyr Pro Pro Leu Pro Ala Gly Tyr Met Thr Lys Glu Leu Glu Tyr Phe Pro 1200 185 1190 1195 Glu Thr Asp Lys Val Trp Ile Glu Ile Gly Glu Thr Glu Gly Lys Phe 1210 Ile Val Asp Ser Val Glu Leu Leu Leu Met Glu Glu <210> 3 <211> 3507 <212> DNA <213> Bacillus thuringiensis <220> <221> CDS <222> (1)..(3507) <400> 3 atg gag ata aat aat cag aac caa tgc ata cca tat aat tgc tta agt Met Glu Ile Asn Asn Gln Asn Gln Cys Ile Pro Tyr Asn Cys Leu Ser aag cct gag gaa gta ttt ttg gat ggg gag agg ata tta cct gat atc 96 Lys Pro Glu Glu Val Phe Leu Asp Gly Glu Arg Ile Leu Pro Asp Ile 20 gat cca ctc gaa gtt tct ttg tcg ctt ttg caa ttt ctt ttg aat aac 144 Asp Pro Leu Glu Val Ser Leu Ser Leu Leu Gln Phe Leu Leu Asn Asn 45 40 35 ttt gtt ccg ggg ggg ggg ttt att tca gga tta att gac aaa ata tgg 192 Phe Val Pro Gly Gly Gly Phe Ile Ser Gly Leu Ile Asp Lys Ile Trp 50 55 ggg gct ttg aga cca tct gaa tgg gaa tta ttt ctt gca cag att gaa Gly Ala Leu Arg Pro Ser Glu Trp Glu Leu Phe Leu Ala Gln Ile Glu 65 cag ttg att gat cga aga ata gaa gca aca gta aga gca aaa gca atc 288 Gln Leu Ile Asp Arg Arg Ile Glu Ala Thr Val Arg Ala Lys Ala Ile 85 336 gct gaa tta gaa ggt tta ggg aga agt tat caa cta tat gga gag gca Ala Glu Leu Glu Gly Leu Gly Arg Ser Tyr Gln Leu Tyr Gly Glu Ala

105

100

ttt Phe	aaa Lys	gag Glu 115	tgg Trp	gaa Glu	aaa Lys	act Thr	cca Pro 120	gat Asp	aac Asn	aca Thr	gcg Ala	gct Ala 125	cgg Arg	tct Ser	aga Arg	384
				ttt Phe												432
cct Pro 145	tcg Ser	ttt Phe	cgg Arg	gtt Val	tcc Ser 150	gga Gly	ttt Phe	gaa Glu	gtg Val	cca Pro 155	ctt Leu	cta Leu	tcg Ser	gtt Val	tat Tyr 160	480
acc Thr	caa Gln	gca Ala	gct Ala	aat Asn 165	ttg Leu	cat His	ctc Leu	gct Ala	cta Leu 170	tta Leu	aga Arg	gat Asp	tct Ser	gtt Val 175	att Ile	528
ttt Phe	gga Gly	gag Glu	aga Arg 180	tgg Trp	gga Gly	ttg Leu	tcg Ser	act Thr 185	aca Thr	aat Asn	gtc Val	aat Asn	gat Asp 190	atc Ile	tat Tyr	576
aat Asn	aga Arg	caa Gln 195	gtt Val	aag Lys	aga Arg	att Ile	cat His 200	gaa Glu	tat Tyr	agc Ser	gat Asp	cat His 205	tgt Cys	gta Val	gat Asp	624
acg Thr	tat Tyr 210	aaa Lys	aca Thr	gaa Glu	tta Leu	gaa Glu 215	cgt Arg	cta Leu	gag Glu	ttt Phe	aga Arg 220	tct Ser	ata Ile	gcg Ala	caa Gln	672
tgg Trp 225	aga Arg	ata Ile	tat Tyr	aat Asn	cag Gln 230	ttt Phe	aga Arg	agg Arg	gaa Glu	ttg Leu 235	aca Thr	cta Leu	acg Thr	gta Val	tta Leu 240	720
				ctt Leu 245												768
				caa Gln												816
				ggt Gly												.864
att Ile	gaa Glu 290	cgg Arg	cag Gln	cta Leu	agg Arg	cga Arg 295	cca Pro	cat His	ctt Leu	atg Met	gac Asp 300	ttc Phe	ttt Phe	aac Asn	tcc Ser	912
atg Met 305	acc Thr	atg Met	tat Tyr	aca Thr	tca Ser 310	gat Asp	aat Asn	aga Arg	cgg Arg	gaa Glu 315	tat Tyr	tat Tyr	tgg Trp	tca Ser	gga Gly 320	960
ctt	gaa	atg	acg	gct	acţ	ctt	act	tca	gga	aat	caa	gtg	tca	ttc	cct	1008

Leu	Glu	Met	Thr	Ala 325	Thr	Leu	Thr	Ser	Gly 330	Asn	Gln	Val	Ser	Phe 335	Pro	
											gta Val					1056
											cca Pro					1104
											380 GJA aaa					1152
											aca Thr					1200
cgt Arg	gga Gly	aca Thr	gta Val	gat Asp 405	tca Ser	tta Leu	gtc Val	agc Ser	ata Ile 410	ccg Pro	cca Pro	caa Gln	gat Asp	tat Tyr 415	agc Ser	1248
gta Val	cca Pro	ccg Pro	cac His 420	agg Arg	gjå aaa	tat Tyr	agt Ser	cat His 425	tta Leu	tta Leu	agt Ser	cac His	gtt Val 430	acg Thr	atg Met	1296
											cgt Arg					1344
aga Arg	aat Asn 450	ata Ile	att Ile	tat Tyr	cca Pro	gat Asp 455	agt Ser	atc Ile	act Thr	caa Gln	atc Ile 460	cca Pro	gta Val	gtt Val	aag Lys	1392
	Ser										gga Gly					1440
ggt Gly	gga Gly	gat Asp	tta Leu	ata Ile 485	agc Ser	cta Leu	cct Pro	gta Val	aat Asn 490	aac Asn	ttt Phe	act Thr	cat His	ttc Phe 495	cga Arg	1488
											cgt Arg					1536
			Asp								gga Gly					1584

											tat Tyr 540				1632
_		_						_	_		caa Gln			_	1680
											tta Leu				1728
											gat Asp				1776
			_	_				_			aat Asn				1824
											aac Asn 620				1872
											gta Val				1920
											aag Lys				1968
											gag Glu				2016
	Asp		Asn		Arg	Gly	Ile	Asn		Glu	caa Gln	Asp	Arg		2064
											aac Asn 700				2112
			_								gcg Ala				2160
											aaa Lys				2208

									agt Ser							2256
ttg Leu	atc Ile	cgt Arg 755	tat Tyr	aat Asn	gca Ala	aaa Lys	tat Tyr 760	gaa Glu	acg Thr	tta Leu	aat Asn	gtg Val 765	cca Pro	ggt Gly	atg Met	2304
									agt Ser							2352
gaa Glu 785	ccg Pro	aat Asn	cga Arg	tgt Cys	gtg Val 790	cca Pro	caa Gln	ctt Leu	gaa Glu	tgg Trp 795	aat Asn	ccc Pro	gat Asp	ttc Phe	gat Asp 800	2400
tgt Cys	tcc Ser	tgc Cys	aga Arg	gac Asp 805	gga Gly	gaa Glu	aaa Lys	tgt Cys	gcg Ala 810	cat His	cat His	tcg Ser	cat His	cat His 815	ttc Phe	2448
tcc Ser	ttg Leu	gac Asp	att Ile 820	gat Asp	gtt Val	gga Gly	tgt Cys	aca Thr 825	gac Asp	ttg Leu	aat Asn	gag Glu	aac Asn 830	cta Leu	ggt Gly	2496
ata Ile	tgg Trp	gtg Val 835	ata Ile	ttc Phe	aag Lys	att Ile	aag Lys 840	aca Thr	cag Gln	gat Asp	ggt Gly	cat His 845	gca Ala	aga Arg	cta Leu	2544
gga Gly	aat Asn 850	cta Leu	gaa Glu	ttt Phe	ctc Leu	gaa Glu 855	gag Glu	aaa Lys	ccg Pro	tta Leu	tta Leu 860	gga Gly	gaa Glu	gcg Ala	tta Leu	2592
gcc Ala 865	cgt Arg	gtg Val	aag Lys	aga Arg	gcg Ala 870	gag Glu	aaa Lys	aaa Lys	tgg Trp	aga Arg 875	gac Asp	aaa Lys	cgt Arg	gaa Glu	ata Ile 880	2640
ttg Leu	caa Gln	tca Ser	gaa Glu	act Thr 885	aat Asn	att Ile	gtt Val	Tyr	aaa Lys 890	Glu	gca Ala	aaa Lys	gaa Glu	gct Ala 895	gta Val	2688
gat Asp	ggt Gly	tta Leu	ttc Phe 900	Val	gat Asp	tcc Ser	caa Gln	tat Tyr 905	gag Glu	aga Arg	tta Leu	caa Gln	tct Ser 910	gat Asp	acg Thr	2736
aat Asn	atc Ile	gcc Ala 915	Met	att Ile	cat His	gcg Ala	gca Ala 920	gat Asp	aaa Lys	cgc Arg	gtt Val	cac His 925	cga Arg	atc Ile	cga Arg	2784
gag Glu	gct Ala 930	tat Tyr	ctg Leu	cca Pro	gag Glu	ctc Leu 935	tcc Ser	gtg Val	att Ile	cca Pro	ggt Gly 940	gtc Val	aat Asn	gca Ala	gcg Ala	2832

att ttc gaa gaa tta gaa gga cgt att ttc aca gcc tac tct cta tat Ile Phe Glu Glu Leu Glu Gly Arg Ile Phe Thr Ala Tyr Ser Leu Tyr 945 950 955 960	2880
gat gcg aga aat gtc att aaa aac ggt gat ttc aat aat ggc tta tca Asp Ala Arg Asn Val Ile Lys Asn Gly Asp Phe Asn Asn Gly Leu Ser 965 970 975	2928
tgc tgg aat gtg aaa ggg cat gta gat ata aaa cag aat ggt cat cgt Cys Trp Asn Val Lys Gly His Val Asp Ile Lys Gln Asn Gly His Arg 980 985 990	2976
tct gtt ctt gtt atc cca gaa tgg gaa gca caa gtt tca caa gaa gtt Ser Val Leu Val Ile Pro Glu Trp Glu Ala Gln Val Ser Gln Glu Val 995 1000 1005	3024
cgt gta tgt cca ggt cgt ggc tac atc ctt cgt gtt aca gcg aac aaa Arg Val Cys Pro Gly Arg Gly Tyr Ile Leu Arg Val Thr Ala Asn Lys 1010 1015 1020	3072
gaa gga tat gga gaa ggc tgc gta acg att cat gag att gag aat cat Glu Gly Tyr Gly Glu Gly Cys Val Thr Ile His Glu Ile Glu Asn His 1025 1030 1035 1040	3120
act gaa aaa ctg aag ttt aga aac tgc gaa gaa gag gaa gta tat cca Thr Glu Lys Leu Lys Phe Arg Asn Cys Glu Glu Glu Glu Val Tyr Pro 1045 1050 1055	3168
aac aat aca gga acg tgt aat gat tat act gca cac caa caa ggt aca Asn Asn Thr Gly Thr Cys Asn Asp Tyr Thr Ala His Gln Gln Gly Thr 1060 1065 1070	3216
gca gga tgc gca gat gca tgt aat acc cgt aat gtc gga tat gaa gat Ala Gly Cys Ala Asp Ala Cys Asn Thr Arg Asn Val Gly Tyr Glu Asp 1075 1080 1085	3264
gca tat gaa atg aat act aca aca tct gtt aat tac aaa ccg act tac Ala Tyr Glu Met Asn Thr Thr Thr Ser Val Asn Tyr Lys Pro Thr Tyr 1090 1095 1100	3312
gag gaa gaa gta tat aca gat gga cga aga gat aat cct tgt gaa atg Glu Glu Glu Val Tyr Thr Asp Gly Arg Arg Asp Asn Pro Cys Glu Met 1105 1110 1115	•
gaa aga ggt tac aca cca tta cca gtt ggt tat gta aca aaa gaa tta Glu Arg Gly Tyr Thr Pro Leu Pro Val Gly Tyr Val Thr Lys Glu Leu 1125 1130 1135	3408
gaa tac ttc cct gaa aca aat aca gta tgg att gaa att gga gaa acg Glu Tyr Phe Pro Glu Thr Asn Thr Val Trp Ile Glu Ile Gly Glu Thr 1140 1145 1150	3456

•

gaa ggg aag ttt att gta gac agt gtc gaa tta ctc ctt atg gaa gaa 3504
Glu Gly Lys Phe Ile Val Asp Ser Val Glu Leu Leu Met Glu Glu
1155 1160 1165
tag

<210> 4

<211> 1168 <212> PRT

<213> Bacillus thuringiensis

<400> 4

Met Glu Ile Asn Asn Gln Asn Gln Cys Ile Pro Tyr Asn Cys Leu Ser

Lys Pro Glu Glu Val Phe Leu Asp Gly Glu Arg Ile Leu Pro Asp Ile 20 25 30

Asp Pro Leu Glu Val Ser Leu Ser Leu Leu Gln Phe Leu Leu Asn Asn 35 40 45

Phe Val Pro Gly Gly Gly Phe Ile Ser Gly Leu Ile Asp Lys Ile Trp 50 55 60

Gly Ala Leu Arg Pro Ser Glu Trp Glu Leu Phe Leu Ala Gln Ile Glu
65 70 75 80

Gln Leu Ile Asp Arg Arg Ile Glu Ala Thr Val Arg Ala Lys Ala Ile 85 90 95

Ala Glu Leu Glu Gly Leu Gly Arg Ser Tyr Gln Leu Tyr Gly Glu Ala 100 105 110

Phe Lys Glu Trp Glu Lys Thr Pro Asp Asn Thr Ala Ala Arg Ser Arg 115 120 125

Val Thr Glu Arg Phe Arg Ile Ile Asp Ala Gln Ile Glu Ala Asn Ile 130 135 140

Pro Ser Phe Arg Val Ser Gly Phe Glu Val Pro Leu Leu Ser Val Tyr 145 150 155 160

Thr Gln Ala Ala Asn Leu His Leu Ala Leu Leu Arg Asp Ser Val Ile 165 170 175

Phe Gly Glu Arg Trp Gly Leu Ser Thr Thr Asn Val Asn Asp Ile Tyr 180 185 190

Asn Arg Gln Val Lys Arg Ile His Glu Tyr Ser Asp His Cys Val Asp 195 200 205

- Thr Tyr Lys Thr Glu Leu Glu Arg Leu Glu Phe Arg Ser Ile Ala Gln 210 215 220
- Trp Arg Ile Tyr Asn Gln Phe Arg Arg Glu Leu Thr Leu Thr Val Leu 225 230 235 240
- Asp Ile Val Ala Leu Phe Pro Asn Tyr Asp Gly Arg Leu Tyr Pro Ile 245 250 255
- Arg Thr Ile Ser Gln Leu Thr Arg Glu Ile Tyr Thr Ser Pro Val Ser 260 265 270
- Glu Phe Tyr Tyr Gly Pro Ile Tyr Asn His Asn Met Ile Gly Thr Phe 275 280 285
- Ile Glu Arg Gln Leu Arg Arg Pro His Leu Met Asp Phe Phe Asn Ser 290 295 300
- Met Thr Met Tyr Thr Ser Asp Asn Arg Arg Glu Tyr Tyr Trp Ser Gly 305 310 315
- Leu Glu Met Thr Ala Thr Leu Thr Ser Gly Asn Gln Val Ser Phe Pro 325 330 335
- Leu Ala Gly Thr Arg Gly Asn Ser Ala Pro Pro Val Ser Val Arg Lys 340 345 350
- Thr Gly Glu Gly Ile Tyr Arg Ile Leu Ser Glu Pro Phe Tyr Ser Ala 355 360 365
- Pro Phe Leu Gly Thr Ser Val Leu Gly Ser Arg Gly Glu Glu Phe Ala 370 375 380
- Phe Ala Ser Asn Thr Thr Thr Ser Leu Pro Ser Thr Ile Tyr Arg Asn 385 390 395 400
- Arg Gly Thr Val Asp Ser Leu Val Ser Ile Pro Pro Gln Asp Tyr Ser 405 410 415
- Val Pro Pro His Arg Gly Tyr Ser His Leu Leu Ser His Val Thr Met 420 425 430
- His Asn Ser Ser Pro Ile Phe His Trp Thr His Arg Ser Ala Thr Pro 435 440 445
- Arg Asn Ile Ile Tyr Pro Asp Ser Ile Thr Gln Ile Pro Val Val Lys 450 455 460
- Ala Ser His Leu Ser Gly Gly Ser Val Ile Lys Gly Pro Gly His Thr 465 470 475 480
- Gly Gly Asp Leu Ile Ser Leu Pro Val Asn Asn Phe Thr His Phe Arg 485 490 495

- Ile Pro Phe Gln Ala Asn Thr Pro Gln Arg Tyr Arg Ile Arg Ile Cys
 500 505 510
- Tyr Ala Ala Asp Ser Asp Gly Thr Leu Asp Ser Gly Val Phe Leu Ser 515 520 525
- Ala Ala Ala Gly Asp Gly Phe Asn Thr Thr Ser Tyr Arg Ala Thr Met 530 535 540
- Ser Pro Glu Gly Ser Leu Thr Ser Arg Asp Phe Gln Phe Leu Asp Leu 545 550 555 560
- Asn Thr Ser Phe Thr Ser Asp Val Ala Ser Asn Leu Trp Leu His Phe 565 570 575
- Ile Arg Tyr Ile Arg Pro Gly Asn Leu Tyr Ile Asp Arg Ala Glu Phe 580 585 590
- Ile Pro Val Asp Ala Thr Phe Glu Ala Gly Tyr Asn Leu Glu Arg Ala
 595 600 605
- Gln Lys Ala Val Asn Ala Leu Phe Thr Ser Thr Asn Gln Lys Gly Leu 610 615 620
- Gln Thr Asp Val Thr Asp Tyr His Ile Asp Gln Val Ser Asn Leu Val 625 630 635 640
- Asp Cys Leu Ser Asp Glu Phe Cys Leu Asp Glu Lys Arg Lys Leu Ser 645 650 655
- Glu Lys Val Lys Gln Ala Lys Arg Leu Ser Asp Glu Arg Asn Leu Leu 660 665 670
- Gln Asp Ser Asn Phe Arg Gly Ile Asn Arg Glu Gln Asp Arg Gly Trp 675 680 685
- Arg Gly Ser Thr Asp Ile Thr Ile Gln Gly Gly Asn Asp Val Phe Lys 690 695 700
- Glu Asn Tyr Val Thr Leu Pro Gly Thr Phe Asp Ala Cys Tyr Pro Thr 705 710 715 720
- Tyr Leu Tyr Gln Lys Val Asp Glu Ser Lys Leu Lys Ala Tyr Thr Arg
 725 730 735
- Tyr Gln Leu Arg Gly Tyr Ile Glu Asp Ser Gln Asp Leu Glu Val Tyr
 740 745 750
- Leu Ile Arg Tyr Asn Ala Lys Tyr Glu Thr Leu Asn Val Pro Gly Met 755 760 765

- Gly Ser Leu Trp Pro Leu Ser Val Glu Ser Pro Ile Gly Lys Cys Gly
 770 775 780
- Glu Pro Asn Arg Cys Val Pro Gln Leu Glu Trp Asn Pro Asp Phe Asp 785 790 795 800
- Cys Ser Cys Arg Asp Gly Glu Lys Cys Ala His His Ser His His Phe 805 810 815
- Ser Leu Asp Ile Asp Val Gly Cys Thr Asp Leu Asn Glu Asn Leu Gly 820 825 830
- Ile Trp Val Ile Phe Lys Ile Lys Thr Gln Asp Gly His Ala Arg Leu 835 840 845
- Gly Asn Leu Glu Phe Leu Glu Glu Lys Pro Leu Leu Gly Glu Ala Leu 850 855 860
- Ala Arg Val Lys Arg Ala Glu Lys Lys Trp Arg Asp Lys Arg Glu Ile 865 870 875 880
- Leu Gln Ser Glu Thr Asn Ile Val Tyr Lys Glu Ala Lys Glu Ala Val 885 890 895
- Asp Gly Leu Phe Val Asp Ser Gln Tyr Glu Arg Leu Gln Ser Asp Thr 900 905 910
- Asn Ile Ala Met Ile His Ala Ala Asp Lys Arg Val His Arg Ile Arg 915 920 925
- Glu Ala Tyr Leu Pro Glu Leu Ser Val Ile Pro Gly Val Asn Ala Ala 930 935 940
- Ile Phe Glu Glu Leu Glu Gly Arg Ile Phe Thr Ala Tyr Ser Leu Tyr 945 950 955 960
- Asp Ala Arg Asn Val Ile Lys Asn Gly Asp Phe Asn Asn Gly Leu Ser 965 970 975
- Cys Trp Asn Val Lys Gly His Val Asp Ile Lys Gln Asn Gly His Arg 980 985 990
- Ser Val Leu Val Ile Pro Glu Trp Glu Ala Gln Val Ser Gln Glu Val 995 1000 1005
- Arg Val Cys Pro Gly Arg Gly Tyr Ile Leu Arg Val Thr Ala Asn Lys 1010 1015 1020
- Glu Gly Tyr Gly Glu Gly Cys Val Thr Ile His Glu Ile Glu Asn His 025 1030 1035 1040
- Thr Glu Lys Leu Lys Phe Arg Asn Cys Glu Glu Glu Glu Val Tyr Pro 1045 1050 1055

1065 1060 Ala Gly Cys Ala Asp Ala Cys Asn Thr Arg Asn Val Gly Tyr Glu Asp 1080 Ala Tyr Glu Met Asn Thr Thr Ser Val Asn Tyr Lys Pro Thr Tyr 1100 1090 1095 Glu Glu Glu Val Tyr Thr Asp Gly Arg Arg Asp Asn Pro Cys Glu Met 1110 1115 Glu Arg Gly Tyr Thr Pro Leu Pro Val Gly Tyr Val Thr Lys Glu Leu 1130 1135 1125 Glu Tyr Phe Pro Glu Thr Asn Thr Val Trp Ile Glu Ile Gly Glu Thr 1150 1145 1140 Glu Gly Lys Phe Ile Val Asp Ser Val Glu Leu Leu Met Glu Glu 1160 1155 <210> 5 <211> 3459 <212> DNA <213> Bacillus thuringiensis <220> <221> CDS <222> (1)..(3459) <400> 5 atg aac cga aat aat caa aat gat tat gaa gtt att gac gct tcc aat 48 Met Asn Arg Asn Asn Gln Asn Asp Tyr Glu Val Ile Asp Ala Ser Asn tgt ggt tgt gcg tca gat gtt gtt caa tac cct ttg gca aga gat 96 Cys Gly Cys Ala Ser Asp Asp Val Val Gln Tyr Pro Leu Ala Arg Asp 25 20 ccg aat gct gta ttc caa aat atg cat tat aaa gat tat ttg caa acg 144 Pro Asn Ala Val Phe Gln Asn Met His Tyr Lys Asp Tyr Leu Gln Thr 45 35 tat gat gga gac tat aca ggt tct ttt ata aat cct aac tta tct att 192 Tyr Asp Gly Asp Tyr Thr Gly Ser Phe Ile Asn Pro Asn Leu Ser Ile 55 50 aat cct aga gat gta ctg caa act gga att aat att gtg gga aga tta Asn Pro Arg Asp Val Leu Gln Thr Gly Ile Asn Ile Val Gly Arg Leu 70 75

Asn Asn Thr Gly Thr Cys Asn Asp Tyr Thr Ala His Gln Gln Gly Thr

cta Leu																288
								cca Pro 105								336
								gag Glu								384
gaa Glu	gca Ala 130	gta Val	gta Val	gly aaa	aca Thr	gca Ala 135	gcg Ala	gat Asp	cat His	tta Leu	acg Thr 140	gga Gly	tta Leu	cac His	gat Asp	432
aat Asn 145	tat Tyr	gag Glu	tta Leu	tat Tyr	gta Val 150	gag Glu	gca Ala	ttg Leu	gaa Glu	gaa Glu 155	tgg Trp	ctg Leu	gaa Glu	aga Arg	ccg Pro 160	480
aat Asn	gct Ala	gct Ala	aga Arg	act Thr 165	aat Asn	cta Leu	ctt Leu	ttt Phe	aat Asn 170	aga Arg	ttt Phe	acc Thr	acc Thr	cta Leu 175	gat Asp	528
agt Ser	ctt Leu	ttt Phe	aca Thr 180	caa Gln	ttt Phe	atg Met	cca Pro	agc Ser 185	ttt Phe	ggt Gly	act Thr	gga Gly	cct Pro 190	gga Gly	agt Ser	576
caa Gln	aac Asn	tac Tyr 195	gca Ala	gtt Val	cca Pro	tta Leu	ctt Leu 200	aca Thr	gta Val	tac Tyr	gca Ala	caa Gln 205	gca Ala	gcg Ala	aac Asn	624
								gct Ala								672
gga Gly 225	ctg Leu	aac Asn	caa Gln	aat Asn	cag Gln 230	Ile	aac Asn	tca Ser	ttc Phe	His	Thr	Arg	caa Gln	caa Gln	gag Glu 240	720
cgt Arg	act Thr	caa Gln	tat Tyr	tat Tyr 245	aca Thr	aat Asn	cat His	tgc Cys	gta Val 250	Thr	acg Thr	tat Tyr	aat Asn	acc Thr 255	ggt Gly	768
tta Leu	gat Asp	aga Arg	tta Leu 260	aga Arg	ggc	aca Thr	aat Asn	act Thr 265	gaa Glu	agt Ser	tgg Trp	tta Leu	aat Asn 270	tat Tyr	cat His	816
cga Arg	ttc Phe	cgt Arg 275	aga Arg	gag Glu	atg Met	aca Thr	tta Leu 280	atg Met	gca Ala	atg Met	gat Asp	tta Leu 285	gtg Val	gcc Ala	tta Leu	864

	cca Pro 290	tac Tyr	tat Tyr	aat Asn	gtg Val	cga Arg 295	caa Gln	tat Tyr	cca Pro	aat Asn	300 Gly Ggg	gca Ala	aat Asn	cca Pro	cag Gln	912
ctt Leu 305	aca Thr	cgt Arg	gaa Glu	ata Ile	tat Tyr 310	acg Thr	gat Asp	cca Pro	atc Ile	gta Val 315	tat Tyr	aat Asn	cca Pro	cca Pro	gct Ala 320	960
aat Asn	cag Gln	gga Gly	atc Ile	tgc Cys 325	cga Arg	cgt Arg	tgg Trp	Gly aaa	aat Asn 330	aat Asn	cct Pro	tat Tyr	aat Asn	aca Thr 335	ttt Phe	1008
tct Ser	gaa Glu	ctt Leu	gaa Glu 340	aat Asn	gct Ala	ttt Phe	att Ile	cgc Arg 345	ccg Pro	cca Pro	cat His	ctt Leu	ttt Phe 350	gat Asp	agg Arg	1056
ttg Leu	aat Asn	aga Arg 355	tta Leu	act Thr	att Ile	tct Ser	aga Arg 360	aac Asn	cga Arg	tat Tyr	aca Thr	gct Ala 365	cca Pro	aca Thr	act Thr	1104
aat Asn	agc Ser 370	tac Tyr	cta Leu	gac Asp	tat Tyr	tgg Trp 375	tca Ser	ggt Gly	cat His	act Thr	tta Leu 380	caa Gln	agc Ser	cag Gln	tat Tyr	1152
gca Ala 385	aat Asn	aac Asn	ccg Pro	acg Thr	aca Thr 390	tat Tyr	gaa Glu	act Thr	agt Ser	tac Tyr 395	ggt Gly	cag Gln	att Ile	acc Tḥr	tct Ser 400	1200
aac Asn	aca Thr	cgt Arg	tta Leu	ttc Phe 405	aat Asn	acg Thr	act Thr	aat Asn	gga Gly 410	gcc Ala	aat Asn	gca Ala	ata Ile	gat Asp 415	tca Ser	1248
agg Arg	gca Ala	aga Arg	aat Asn 420	ttt Phe	ggt Gly	aac Asn	tta Leu	tac Tyr 425	gct Ala	aat Asn	ttg Leu	tat Tyr	ggt Gly 430	gtt Val	agc Ser	1296
tat Tyr	Leu	aat Asn 435	Ile	ttc Phe	cca Pro	aca Thr	ggt Gly 440	Val	atg Met	agt Ser	gaa Glu	atc Ile 445	acc Thr	tca Ser	gcc Ala	1344
cct Pro	aat Asn 450	Thr	tgt Cys	tgg Trp	caa Gln	gac Asp 455	Leu	act Thr	aca Thr	act Thr	gag Glu 460	Glu	cta Leu	cca Pro	cta Leu	1392
gtg Val 465	Asn	aat Asn	aat Asn	ttt Phe	aat Asn 470	Leu	tta Leu	tct Ser	cat His	gtt Val 475	Thr	ttc Phe	tta Leu	cgc Arg	ttc Phe 480	1440
aat Asn	act Thr	act Thr	cag Gln	ggt Gly 485	Gly	ccc Pro	ctt Leu	gca Ala	act Thr 490	· Val	gly ggg	ttt Phe	gta Val	Pro 495	aca Thr	1488

										aat Asn						1536
										gca Ala						1584
ggt Gly	gct Ala 530	act Thr	gtc Val	gtg Val	aaa Lys	ggt Gly 535	cca Pro	gga Gly	ttc Phe	aca Thr	gga Gly 540	gga Gly	gat Asp	gta Val	atc Ile	1632
cga Arg 545	aga Arg	aca Thr	aat Asn	act Thr	ggt Gly 550	gga Gly	ttc Phe	gga Gly	gca Ala	ata Ile 555	agg Arg	gtg Val	tcg Ser	gtc Val	act Thr 560	1680
gga Gly	ccg Pro	cta Leu	Thr	caa Gln 565	cga Arg	tat Tyr	cgc Arg	ata Ile	agg Arg 570	ttc Phe	cgt Arg	tat Tyr	gct Ala	tcg Ser 575	aca Thr	1728
										gga Gly						1776
										cag Gln						1824
tcc Ser	tat Tyr 610	cgt Arg	act Thr	gta Val	gag Glu	ttt Phe 615	aca Thr	act Thr	cct Pro	ttt Phe	aac Asn 620	ttt Phe	aca Thr	caa Gln	agt Ser	1872
caa Gln 625	gat Asp	ata Ile	att Ile	cga Arg	aca Thr 630	tct Ser	atc Ile	cag Gln	gga Gly	ctt Leu 635	agt Ser	gga Gly	aat Asn	Gly 999	gaa Glu 640	1920
gta Val	tac Tyr	Leu	Asp	aga Arg 645	Ile	gaa Glu	atc Ile	atc Ile	cct Pro 650	gta Val	aat Asn	cca Pro	aca Thr	cga Arg 655	gaa Glu	1968
gcg Ala	gaa Glu	gag Glu	gat Asp 660	tta Leu	gaa Glu	gcg Ala	gcg Ala	aag Lys 665	aaa Lys	gcg Ala	gtg Val	gcg Ala	agc Ser 670	ttg Leu	ttt Phe	2016
aca Thr	cgc Arg	aca Thr 675	agg Arg	gac Asp	gga Gly	tta Leu	caa Gln 680	gta Val	aat Asn	gtg Val	aca Thr	gat Asp 685	tat Tyr	caa Gln	gtc Val	2064
gat Asp	caa Gln 690	Ala	gca Ala	aat Asn	tta Leu	gtg Val 695	tca Ser	tgc Cys	tta Leu	tca Ser	gat Asp 700	Glu	caa Gln	tat Tyr	gcg Ala	2112

. .

cat His 705	gat Asp	aaa Lys	aag Lys	atg Met	tta Leu 710	ttg Leu	gaa Glu	gcg Ala	gta Val	cgc Arg 715	gcg Ala	gca Ala	aaa Lys	cgc Arg	ctc Leu 720	2160
agc Ser	cga Arg	gaa Glu	cgt Arg	aac Asn 725	tta Leu	ctt Leu	cag Gln	gat Asp	cca Pro 730	gat Asp	ttt Phe	aat Asn	aca Thr	atc Ile 735	aat Asn	2208
agt Ser	aca Thr	gaa Glu	gaa Glu 740	aat Asn	gga Gly	tgg Trp	aaa Lys	gca Ala 745	agt Ser	aac Asn	ggc Gly	gtt Val	act Thr 750	att Ile	agt Ser	2256
gag Glu	ggc	ggt Gly 755	cca Pro	ttc Phe	tat Tyr	aaa Lys	ggc Gly 760	cgt Arg	gca Ala	ctt Leu	cag Gln	cta Leu 765	gca Ala	agt Ser	gca Ala	2304
cga Arg	gaa Glu 770	aat Asn	tac Tyr	cca Pro	aca Thr	tac Tyr 775	atc Ile	tat Tyr	caa Gln	aaa Lys	gta Val 780	gat Asp	gca Ala	tcg Ser	gag Glu	2352
tta Leu 785	aaa Lys	cct Pro	tat Tyr	aca Thr	cga Arg 790	tat Tyr	aga Arg	ctg Leu	gat Asp	ggg Gly 795	ttc Phe	gtg Val	aag Lys	agt Ser	agt Ser 800	2400
caa Gln	gat Asp	tta Leu	gaa Glu	att Ile 805	gat Asp	ctc Leu	att Ile	cac His	cat His 810	cat His	aaa Lys	gtc Val	cat His	ctt Leu 815	gtg Val	2448
aaa Lys	aat Asn	gta Val	cta Leu 820	gat Asp	aat Asn	tta Leu	gta Val	tct Ser 825	gat Asp	act Thr	tac Tyr	cca Pro	gat Asp 830	gat Asp	tct Ser	2496
tgt Cys	agt Ser	gga Gly 835	atc Ile	aat Asn	cga Arg	tgt Cys	gag Glu 840	gaa Glu	caa Gln	cag Gln	atg Met	gta Val 845	aat Asn	gcg Ala	caa Gln	2544
ctg Leu	gaa Glu 850	Thr	gaa Glu	cat His	cat His	cat His 855	Pro	atg Met	gat Asp	tgc Cys	tgt Cys 860	Glu	gca Ala	gct Ala	caa Gln	2592
aca Thr 865	His	gag Glu	ttt Phe	tct Ser	tcc Ser 870	Tyr	att Ile	gat Asp	aca Thr	999 Gly 875	Asp	tta Leu	aat Asn	tcg Ser	act Thr 880	2640
gta Val	gac	cag Gln	gga Gly	atc Ile 885	Trp	gtg Val	atc Ile	ttt Phe	aaa Lys 890	Val	cga Arg	aca Thr	aca Thr	gat Asp 895	ggt	2688
tat Tyr	gcg Ala	acg Thr	cta Leu 900	Gly	aat Asn	ctt Leu	gaa Glu	ttg Leu 905	Val	gag Glu	gto Val	gga Gly	ccg Pro 910	Leu	ttg Leu	2736

ggt gaa cct Gly Glu Pro 915	cta gaa cgt g Leu Glu Arg G	gaa caa aga Slu Gln Arg 920	gaa aat gcg Glu Asn Ala	aaa tgg aat Lys Trp Asn 925	gca 2784 Ala
gag tta gga Glu Leu Gly 930	aga aaa cgt g Arg Lys Arg A	gca gaa aca Ala Glu Thr 935	gat cgc gtg Asp Arg Val 940	tat caa gat Tyr Gln Asp	gcc 2832 Ala
	atc aat cat t Ile Asn His I 950				
aat cca caa Asn Pro Gln	ata ggg atg g Ile Gly Met <i>I</i> 965	gca gat att Ala Asp Ile	atg gac gct Met Asp Ala 970	caa aat ctt Gln Asn Leu 975	gtc 2928 Val
gca tca att Ala Ser Ile	tca gat gta t Ser Asp Val 1 980	tat agc gat Tyr Ser Asp 985	gca gta ctg Ala Val Leu	caa atc cct Gln Ile Pro 990	gga 2976 Gly
att aac tat Ile Asn Tyr 995	gag att tac a	aca gag ctg Thr Glu Leu 1000	Ser Asn Arg	tta caa caa Leu Gln Gln L005	gca 3024 Ala
tcg tat ctg Ser Tyr Leu 1010	tat acg tct of Tyr Thr Ser A	cga aat gcg Arg Asn Ala 015	gtg caa aat Val Gln Asn 1020	ggg gac ttt Gly Asp Phe	aac 3072 Asn
aac ggg cta Asn Gly Leu 1025	gat agc tgg a Asp Ser Trp 1	aat gca aca Asn Ala Thr	gcg ggt gca Ala Gly Ala 1035	tcg gta caa Ser Val Gln	cag 3120 Gln 1040
gat ggc aat Asp Gly Asn	acg cat ttc f Thr His Phe 1 1045	Leu Val Leu	tct cat tgg Ser His Trp L050	gat gca caa Asp Ala Gln 1055	Val
Ser Gln Gln	ttt aga gtg (Phe Arg Val (1060	cag ccg aat Gln Pro Asn 1065	Cys Lys Tyr	gta tta cgt Val Leu Arg 1070	gta 3216 Val
aca gca gag Thr Ala Glu 1075	aaa gta ggc	ggc gga gac Gly Gly Asp 1080	Gly Tyr Val	act atc cgg Thr Ile Arg 1085	gat 3264 Asp
ggt gct cat Gly Ala His 1090	cat aca gaa His Thr Glu	acg ctt aca Thr Leu Thr 095	ttt aat gca Phe Asn Ala 1100	tgt gat tat Cys Asp Tyr	gat 3312 Asp
ata aat ggc Ile Asn Gly 1105	acg tac gtg Thr Tyr Val	act gat aat Thr Asp Asn	acg tat cta Thr Tyr Leu 1115	aca aaa gaa Thr Lys Glu	gtg 3360 Val 1120

ata ttc tat tca cat aca gaa cac atg tgg gta gag gta aat gaa aca Ile Phe Tyr Ser His Thr Glu His Met Trp Val Glu Val Asn Glu Thr 1130 gaa ggt gca ttt cat ata gat agt att gaa ttc gtt gaa aca gaa aag Glu Gly Ala Phe His Ile Asp Ser Ile Glu Phe Val Glu Thr Glu Lys 1140 1145 taa <210> 6 <211> 1152 <212> PRT <213> Bacillus thuringiensis <400> 6 Met Asn Arg Asn Asn Gln Asn Asp Tyr Glu Val Ile Asp Ala Ser Asn Cys Gly Cys Ala Ser Asp Val Val Gln Tyr Pro Leu Ala Arg Asp Pro Asn Ala Val Phe Gln Asn Met His Tyr Lys Asp Tyr Leu Gln Thr 35 Tyr Asp Gly Asp Tyr Thr Gly Ser Phe Ile Asn Pro Asn Leu Ser Ile Asn Pro Arg Asp Val Leu Gln Thr Gly Ile Asn Ile Val Gly Arg Leu 75 70 Leu Gly Phe Leu Gly Val Pro Phe Ala Gly Gln Leu Val Thr Phe Tyr 90 85 Thr Phe Leu Leu Asn Gln Leu Trp Pro Thr Asn Asp Asn Ala Val Trp 105 Glu Ala Phe Met Ala Gln Ile Glu Glu Leu Ile Asn Gln Arg Ile Ser 120 115 Glu Ala Val Val Gly Thr Ala Ala Asp His Leu Thr Gly Leu His Asp 135 130 Asn Tyr Glu Leu Tyr Val Glu Ala Leu Glu Glu Trp Leu Glu Arg Pro 155 150 Asn Ala Ala Arg Thr Asn Leu Leu Phe Asn Arg Phe Thr Thr Leu Asp 175 165

3459

Ser Leu Phe Thr Gln Phe Met Pro Ser Phe Gly Thr Gly Pro Gly Ser 185

180

- Gln Asn Tyr Ala Val Pro Leu Leu Thr Val Tyr Ala Gln Ala Ala Asn 195 200 205
- Leu His Leu Leu Leu Leu Lys Asp Ala Glu Ile Tyr Gly Ala Arg Trp 210 215 220
- Gly Leu Asn Gln Asn Gln Ile Asn Ser Phe His Thr Arg Gln Gln Glu 225 230 235 240
- Arg Thr Gln Tyr Tyr Thr Asn His Cys Val Thr Thr Tyr Asn Thr Gly 245 250 255
- Leu Asp Arg Leu Arg Gly Thr Asn Thr Glu Ser Trp Leu Asn Tyr His 260 265 270
- Arg Phe Arg Arg Glu Met Thr Leu Met Ala Met Asp Leu Val Ala Leu 275 280 285
- Phe Pro Tyr Tyr Asn Val Arg Gln Tyr Pro Asn Gly Ala Asn Pro Gln 290 295 300
- Leu Thr Arg Glu Ile Tyr Thr Asp Pro Ile Val Tyr Asn Pro Pro Ala 305 310 315 320
- Asn Gln Gly Ile Cys Arg Arg Trp Gly Asn Asn Pro Tyr Asn Thr Phe 325 330 335
- Ser Glu Leu Glu Asn Ala Phe Ile Arg Pro Pro His Leu Phe Asp Arg 340 345 350
- Leu Asn Arg Leu Thr Ile Ser Arg Asn Arg Tyr Thr Ala Pro Thr Thr 355 360 365
- Asn Ser Tyr Leu Asp Tyr Trp Ser Gly His Thr Leu Gln Ser Gln Tyr 370 375 380
- Ala Asn Asn Pro Thr Thr Tyr Glu Thr Ser Tyr Gly Gln Ile Thr Ser 385 390 395 400
- Asn Thr Arg Leu Phe Asn Thr Thr Asn Gly Ala Asn Ala Ile Asp Ser 405 410 415
- Arg Ala Arg Asn Phe Gly Asn Leu Tyr Ala Asn Leu Tyr Gly Val Ser 420 425 430
- Tyr Leu Asn Ile Phe Pro Thr Gly Val Met Ser Glu Ile Thr Ser Ala 435 440 445
- Pro Asn Thr Cys Trp Gln Asp Leu Thr Thr Thr Glu Glu Leu Pro Leu 450 455 460

- Val Asn Asn Asn Phe Asn Leu Leu Ser His Val Thr Phe Leu Arg Phe 465

 Asn Thr Thr Gln Gly Gly Pro Leu Ala Thr Val Gly Phe Val Pro Thr 485
- Tyr Val Trp Thr Arg Gln Asp Val Asp Phe Asn Asn Ile Ile Thr Pro
 500 505 510
- Asn Arg Ile Thr Gln Ile Pro Val Val Lys Ala Tyr Glu Leu Ser Ser 515 520 525
- Gly Ala Thr Val Val Lys Gly Pro Gly Phe Thr Gly Gly Asp Val Ile 530 535 540
- Arg Arg Thr Asn Thr Gly Gly Phe Gly Ala Ile Arg Val Ser Val Thr 545 550 555 560
- Gly Pro Leu Thr Gln Arg Tyr Arg Ile Arg Phe Arg Tyr Ala Ser Thr 565 570 575
- Ile Asp Phe Asp Phe Phe Val Thr Arg Gly Gly Thr Thr Ile Asn Asn 580 585 590
- Phe Arg Phe Thr Arg Thr Met Asn Arg Gly Gln Glu Ser Arg Tyr Glu 595 600 605
- Ser Tyr Arg Thr Val Glu Phe Thr Thr Pro Phe Asn Phe Thr Gln Ser
- Gln Asp Ile Ile Arg Thr Ser Ile Gln Gly Leu Ser Gly Asn Gly Glu 625 630 635 640
- Val Tyr Leu Asp Arg Ile Glu Ile Ile Pro Val Asn Pro Thr Arg Glu 645 650 655
- Ala Glu Glu Asp Leu Glu Ala Ala Lys Lys Ala Val Ala Ser Leu Phe 660 665 670
- Thr Arg Thr Arg Asp Gly Leu Gln Val Asn Val Thr Asp Tyr Gln Val 675 680 685
- Asp Gln Ala Ala Asn Leu Val Ser Cys Leu Ser Asp Glu Gln Tyr Ala 690 695 700
- His Asp Lys Lys Met Leu Leu Glu Ala Val Arg Ala Ala Lys Arg Leu 705 710 715 720
- Ser Arg Glu Arg Asn Leu Leu Gln Asp Pro Asp Phe Asn Thr Ile Asn 725 730 735
- Ser Thr Glu Glu Asn Gly Trp Lys Ala Ser Asn Gly Val Thr Ile Ser 740 745 750

- Glu Gly Gly Pro Phe Tyr Lys Gly Arg Ala Leu Gln Leu Ala Ser Ala 755 760 765
- Arg Glu Asn Tyr Pro Thr Tyr Ile Tyr Gln Lys Val Asp Ala Ser Glu
 770 775 780
- Leu Lys Pro Tyr Thr Arg Tyr Arg Leu Asp Gly Phe Val Lys Ser Ser 785 790 795 800
- Gln Asp Leu Glu Ile Asp Leu Ile His His His Lys Val His Leu Val 805 810 815
- Lys Asn Val Leu Asp Asn Leu Val Ser Asp Thr Tyr Pro Asp Asp Ser 820 825 830
- Cys Ser Gly Ile Asn Arg Cys Glu Glu Gln Gln Met Val Asn Ala Gln 835 840 845
- Leu Glu Thr Glu His His Pro Met Asp Cys Cys Glu Ala Ala Gln 850 855 860
- Thr His Glu Phe Ser Ser Tyr Ile Asp Thr Gly Asp Leu Asn Ser Thr 865 870 875 880
- Val Asp Gln Gly Ile Trp Val Ile Phe Lys Val Arg Thr Thr Asp Gly 885 890 895
- Tyr Ala Thr Leu Gly Asn Leu Glu Leu Val Glu Val Gly Pro Leu Leu 900 905 910
- Gly Glu Pro Leu Glu Arg Glu Gln Arg Glu Asn Ala Lys Trp Asn Ala 915 920 925
- Glu Leu Gly Arg Lys Arg Ala Glu Thr Asp Arg Val Tyr Gln Asp Ala 930 935 940
- Lys Gln Ser Ile Asn His Leu Phe Val Asp Tyr Gln Asp Gln Gln Leu 945 950 955 960
- Asn Pro Gln Ile Gly Met Ala Asp Ile Met Asp Ala Gln Asn Leu Val 965 970 975
- Ala Ser Ile Ser Asp Val Tyr Ser Asp Ala Val Leu Gln Ile Pro Gly
 980 985 990
- Ile Asn Tyr Glu Ile Tyr Thr Glu Leu Ser Asn Arg Leu Gln Gln Ala 995 1000 1005
- Ser Tyr Leu Tyr Thr Ser Arg Asn Ala Val Gln Asn Gly Asp Phe Asn 1010 1015 1020

Asn Gly Leu Asp Ser Trp Asn Ala Thr Ala Gly Ala Ser Val Gln Gln 1035 1030 025 Asp Gly Asn Thr His Phe Leu Val Leu Ser His Trp Asp Ala Gln Val 1050 1045 Ser Gln Gln Phe Arg Val Gln Pro Asn Cys Lys Tyr Val Leu Arg Val 1060 1065 Thr Ala Glu Lys Val Gly Gly Gly Asp Gly Tyr Val Thr Ile Arg Asp 1080 1075 Gly Ala His His Thr Glu Thr Leu Thr Phe Asn Ala Cys Asp Tyr Asp 1100 1095 Ile Asn Gly Thr Tyr Val Thr Asp Asn Thr Tyr Leu Thr Lys Glu Val 1120 1115 1110 105 Ile Phe Tyr Ser His Thr Glu His Met Trp Val Glu Val Asn Glu Thr 1125 1130 Glu Gly Ala Phe His Ile Asp Ser Ile Glu Phe Val Glu Thr Glu Lys 1145 1140 <210> 7 <211> 20 <212> DNA <213> Artificial Sequence <223> Description of Artificial Sequence: primer cry1B.fw <400> 7 20 cagtccaaac gggtataaac <210> 8 <211> 20 <212> DNA <213> Artificial Sequence <223> Description of Artificial Sequence: primer B.R <400> 8 20 ctgcttcgaa ggttgcagta <210> 9 <211> 20

<212> DNA

```
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer B.F
<400> 9
                                                                    20
cagcgtatta agtcgatgga
<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer JFW
<400> 10
                                                                    20
gcagctaatg ctaccacatc
<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer JRV
<400> 11
                                                                    20
gtggcggtat gctgactaat
<210> 12
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer 9FW
<400> 12
                                                                    20
gyttttattc gcccgccaca
<210> 13
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> Description of Artificial Sequence: primer 9RV
```

<400> 13 cgacagtags acccactact